

**A COMPARATIVE EVALUATION OF ANTIMICROBIAL
EFFICACY OF CINNAMON AND GARLIC AS ENDODONTIC
IRRIGANTS AGAINST ENTEROCOCCUS FAECALIS-
AN IN VITRO STUDY**

Dissertation submitted to

The Tamil Nadu Dr. M G R Medical University

In the partial fulfillment of the degree of

MASTER OF DENTAL SURGERY



Branch IV

CONSERVATIVE DENTISTRY AND ENDODONTICS

2010-2013

CERTIFICATE

This is to certify that this dissertation titled "A comparative evaluation of antimicrobial efficacy Of Cinnamon And Garlic as endodontic irrigants against Enterococcus faecalis-An invitro study", is a bonafide record of the work done by Dr.Santhini Gopalakrishnan Nair under our guidance during his/her post graduate study during the period of 2010-2013 under THE TAMIL NADU Dr.MGR MEDICAL UNIVERSITY, CHENNAI, inpartial fulfilment for the degree of MASTER OF DENTAL SURGERY IN CONSERVATIVE DENTISTRY & ENDODONTICS, BRANCH IV. It has not been submitted (partial or full) for the award of any other degree or diploma.

Dr.Rajesh.S M.D.S

Professor, HOD and Guide

Department of Conservative Dentistry &Endodontics



Dr. S. Rajesh MDS.
Professor & Head
Dept. of Conservative Dentistry and Endodontics
Sree Mookambika Institute of Dental Sciences
Kulashekaram



ACKNOWLEDGEMENT

I take this opportunity to thank my teacher and mentor **Dr S Rajesh MDS**, *Professor, Guide and Head of the Department, Department of Conservative Dentistry and Endodontics, Sree Mookambika Institute of Dental Sciences, Kulasekharam, Kanyakumari District*, for his help, support and patient guidance during this study and also right from the time I joined for this course. His immense knowledge and remarkable insight into the subject of Conservative Dentistry and Endodontics has made the topic easier to understand and remember.

I would also like to acknowledge and thank **Dr Vijay Mathai, MDS** *Professor, Department of Conservative Dentistry and Endodontics, Sree Mookambika Institute of Dental Sciences, Kulasekharam*, for being a source of inspiration and motivation in continuing the study.

I am deeply indebted to **Dr K C Mariamma MDS**, *Reader, Department of Conservative Dentistry and Endodontics, Sree Mookambika Institute of Dental Sciences, Kulasekharam*, for her timely help, support and encouragement throughout the study.

My heartfelt and sincere thanks to **Dr Mano Christaine MDS**, *Reader*. His nature of allowing a great deal of freedom in work and adhering strictly at the same time to the guiding principles and high teaching standards has been a source of encouragement to me and beneficial to my professional improvement as a post-graduate student.

Dr Jothish Ravi MDS, *Senior Lecturer*. Without his help and guidance during the project, this study would not have been possible. I also thank him for making

Conservative Dentistry and Endodontics an interesting subject, in his own unique style. and **Dr. T.S. Manoj Kumar BDS** for their valuable help and guidance.

I am grateful to **V.Gopalakrishnan**, *Head of the Department of Microbiology, Pankajakasthuri Research centre, Trivandrum* for patiently explaining and helping me to understand and perform the experiment for the study.

I am thankful to **Mr.Sharath Babu** for providing me with his timely statistical analysis involved in this study.

I gratefully acknowledge my batch mate **Dr Manu Unnikrishnan** and my fellow post graduates, **Dr Suji Susan, Dr Rajalakshmy, Dr Sarah Christopher and Dr Rahul S** for their motivation and encouraging words.

I would like to thank all my family members for their prayers, constant enthusiasm and immense support.

The small endeavour of mine would not have been possible without the help and effort from Print Land Printers, Nagercoil. I thank them for their sincere support.

Above all I thank the Almighty God for walking with me in each and every step of this study and making it a successful one.

CONTENTS

SL NO:	INDEX	PAGE NO:
1.	List of Abbreviations	I
2.	List of Tables	II
3.	List of Graphs	III
4.	List of Figures	IV
5.	Abstract	V-VII
6.	Introduction	1-5
7.	Aims & Objectives	6
8.	Review of Literature	7-32
9.	Materials & Methods	33-39
10.	Results and Observations	40-42
11.	Discussion	43-49
12.	Summary & Conclusion	50-51
13.	Tables	VIII-XI
14.	Graphs	XII-XIII
15.	Figures	XIV-XXIII
16.	Bibliography	XXIV-XLI

LIST OF ABBREVIATIONS

E.faecalis	-	Enterococcus Faecalis
Ca (OH) ₂	-	Calcium Hydroxide
NaOCl	-	Sodium Hypochlorite
CHX	-	Chlorhexidinedigluconate
EDTA	-	Ethylene di-amine tetra acetic acid
MTAD	-	Mixture of tetracycline, acid and detergent
µm	-	Micro meter
ANOVA	-	Analysis of Variance
SEM	-	Scanning Electron Microscopy
ATCC	-	American Type Culture Collection
MBC	-	Minimum bactericidal concentration
MIC	-	Minimum inhibitory concentration
CFU	-	Colony forming unit
BHI	-	Brain heart infusion broth
SPSS	-	Statistical Package for Social Sciences

LIST OF TABLES

Table 1	Mean values of number of viable colonies of bacteria in each group at 10^{-1} dilution
Table 2	Mean values of number of viable colonies of bacteria in each group at 10^{-3} dilution
Table 3	Mean values of number of viable colonies of bacteria in each group at 10^{-6} dilution
Table 4	Multiple comparison of mean number of viable colonies of bacteria in the different groups
Table 5	Mean values of number and percentage of samples with growth of bacteria in each group at 10^{-1} dilution
Table 6	Mean values of number and percentage of samples with growth of bacteria in each group at 10^{-3} dilution
Table 7	Mean values of number and percentage of samples with growth of bacteria in each group at 10^{-6} dilution
Table 8	Multiple comparisons of mean values of number and percentage of samples with growth of bacteria between the groups.

LIST OF GRAPHS

Graph 1	Mean values of number of viable colonies of bacteria in each group at 10^{-1} dilution
Graph2	Multiple comparison of mean number of viable colonies of bacteria in the different groups
Graph3	Multiple comparisons of mean values of number and percentage of samples with growth of bacteria.

LIST OF FIGURES

1	Specimens stored in normal saline after biomechanical preparation
2	Specimens immersed in streptococcus selection broth for autoclaving
3	Specimens subjected to autoclaving
4	Laminar flow
5	Inoculation of specimens with E.faecalis suspension using tuberculin syringes
6	Specimens kept for incubation
7	SEM picture showing E.faecalis colonies
8	Irrigation of specimens with the test irrigants
9	Sample preparation using H files.
10	Collection of dentinal shavings in test tube containing normal saline
11	Agar plates showing complete inhibition by 5.25% NaOCl
12	Agar plate showing complete inhibition by 2% CHX
13	Agar plates (Garlic) showing a) complete inhibition b) presence of CFU
14	Agar plates (Cinnamon) showing a) complete inhibition b) presence of CFU
15	Agar plate (Saline) showing innumerable CFU.

Abstract

A COMPARATIVE EVALUATION OF ANTIMICROBIAL EFFICACY OF CINNAMON AND GARLIC AS ENDODONTIC IRRIGANTS AGAINST ENTEROCOCCUS FAECALIS – AN IN VITRO STUDY

ABSTRACT

Introduction

Eliminating microorganism from root canal system is possible only by a thorough chemomechanical preparation which is accomplished by proper instrumentation along with irrigants and intracanal medicaments. However complete sterilization of pulp space is not always achieved due to extremely complex anatomy. Persistent endodontic infections are mainly due to retention of microorganisms in the dentinal tubules. Enterococcus faecalis and Candida albicans constitute 77% of the persistent asymptomatic infections. It is important to appreciate that while hand and rotary instruments produce shape, it is the irrigants that clean the root canal system. Irrigants not only are important for the removal of debris and dentinal chips produced during cleaning and shaping, but are of clinical importance in the eradication of the radicular infection.

Aims & Objectives:

To evaluate the antimicrobial efficacy of Cinnamon and Garlic as endodontic irrigants against Enterococcus faecalis with comparison of 5.25% NaOCl and 2% CHX.

Methodology

Fifty freshly extracted intact human mandibular premolars were decoronated and biomechanically prepared using 15-50 K files. The specimens were stored in normal saline until autoclaving. Specimens were autoclaved in separate steel containers containing BHI broth. The specimens were inoculated with *E. faecalis* suspension and incubated for 21 days in an incubator. The broth was replaced on alternate days. After 21 days the specimens were retrieved and divided into five groups (N=10). One tooth was subjected for SEM evaluation to confirm the penetration of bacteria. Five groups were treated with the respective irrigating solutions for 5 minutes. The sample preparation was done using H files and transferred into test tubes containing 10 ml sterile normal saline (10^{-1}). Three dilutions 10^{-1} , 10^{-3} and 10^{-6} were used for the count. From this one ml from each dilution was pipetted on to a sterile 100 mm diameter in duplicate. To each of these plates 15 ml of agar medium, melted and cooled to 45°C , was added, mixed well and allowed to solidify. These plates were incubated for two days at 37°C . After incubation the number of colonies was counted in suitable plates. The number of the colonies multiplied by the dilution factor gives the total number of CFU in the scrapings per tooth.

Results

CFU was determined in three dilutions 10^{-1} , 10^{-3} , 10^{-6} for all the five groups. 5.25% NaOCl and 2% CHX showed complete inhibition in all the samples. Garlic and Cinnamon showed reduction in CFU in all the samples tested in 10^{-1} dilution. Complete inhibition was observed in 2 samples of Garlic and 1 sample of Cinnamon. Saline showed innumerable colonies in the same dilution.

Summary

The present study was done to evaluate the antimicrobial efficacy of Cinnamon and Garlic as endodontic irrigants against *E.faecalis* when compared to NaOCl and CHX. The study concluded that 5.25% NaOCl and 2% CHX showed complete inhibition where as Cinnamon and Garlic showed inhibition which suggest that they have an antimicrobial action but not up to the extent of NaOCl and CHX.

Clinical Implications

The use of herbal alternatives as a root canal irrigant might prove to be advantageous considering the several undesirable characteristics of standard irrigants and frequently used antimicrobials. Further research is warranted to conclusively recommend herbal solutions as a root canal irrigant.

Introduction

Successful endodontic treatment depends on healthy periradicular tissue. The importance of bacteria in pulpal pathosis has been demonstrated in the past (1). Primary endodontic infections are polymicrobial and are dominated by obligatory anaerobic bacteria (2). Eliminating microorganism from root canal system is possible only by a thorough chemomechanical preparation (3) which is accomplished by proper instrumentation along with irrigants and intracanal medicaments (4). However complete sterilization of pulp space is not always achieved due to extremely complex anatomy (5). Even after meticulous chemomechanical preparation bacteria can still be recovered from canals (6). Persistent endodontic infections are mainly due to retention of microorganism in the dentinal tubules. *Enterococcus faecalis* is the primary organism detected in persistent asymptomatic infections (7). *Enterococcus faecalis* is a facultative anaerobic gram positive rod which can invade the dentinal tubules (5) endure prolonged periods of starvation and possess certain virulence factors and lytic enzymes (8,9). Its mode of action is through biofilm formation which helps it resist destruction by enabling the bacteria to become 1000 times more resistant to most commonly used irrigants and intracanal medicaments(8).

Thorough chemomechanical preparation plays a key role in the success of endodontic treatment. It is important to appreciate that while hand and rotary instruments produce shape, it is the irrigants that clean the root canal system. Irrigants not only are important for the removal of debris and dentinal chips produced during cleaning and shaping, but are of clinical importance in the eradication of the radicular infection (10).

An ideal irrigant should have (11, 12)

1. Broad antimicrobial spectrum

2. Mechanically flushes out the debris
3. Nontoxic and biocompatible
4. Dissolves necrotic and vital pulp tissue
5. Serves as a lubricant
6. Removes the smear layer
7. Low surface tension

NaOCl is a widely used irrigant as it covers most of the requirements for endodontic irrigants than any other known compound. Hypochlorite has the unique capacity to dissolve the necrotic tissue and the organic components of smear layer. Studies conducted so far point out to evidence that is strongly in favour of NaOCl as the main irrigant (11,12). But it has some disadvantages such as high toxicity, unpleasant taste, corrosive to instruments, inability to remove the inorganic portion of smear layer and reduction in elastic modulus and flexural strength of dentin (13). The difference in cell structure, thickness and endotoxin formation of facultative gram positive anaerobes makes it resistant to low concentration of NaOCl when compared to gram negative organisms. Increasing the concentration has its consequences. The purpose of introducing 17% EDTA into the irrigation regimen was to remove the smear layer which will facilitate penetration of irrigant to greater depth of the dentinal tubule.

Another widely accepted irrigant is 2% Chlorhexidine digluconate. It has a broad spectrum antimicrobial action, low toxicity and property of substantivity, but it cannot dissolve the organic substrate and necrotic tissue from the root canal system (11,12). Allergic reactions have also been reported against 2% CHX such as contact

dermatitis, desquamative gingivitis, discolouration of the teeth and tongue and dysgeusia (12). Due to the potential risk of adverse effects of systemic applications, and the ineffectiveness of systemic antibiotics in necrotic or pulpless teeth and the periapical tissues, the local application of antibiotics emerged as an effective mode of delivering antibiotics to infected root canals. Some of the antibiotic based irrigants are MTAD and Tetraclean (14). Other irrigants recently introduced are HEBP, EDTA-T, Chlorine dioxide, Silver diamine fluoride, Triclosan and Gantrez, Ozonated water, Photon activated disinfection (10). The constant increase in antimicrobial resistance and side effects caused by synthetic drugs has prompted researchers to look for herbal alternatives. Scientists have realized an immense potential in natural products made from medicinal plants to serve as alternate source of combating infections in human beings (15,16).

Herbs in dentistry have become more popular due to easy availability, cost effectiveness, low toxicity and lack of microbial resistance and increased shelf life (13). The changing trends from conventional irrigants to herbal extracts began in 2003 when propolis was compared with saline and NaOCl as root canal irrigants (17). Later Morinda citrifolia juice was used in conjunction with EDTA as a possible alternative to NaOCl (17). Other herbal alternatives such as Triphala, Green tea polyphenols were also recently evaluated as endodontic irrigants (16).

Spices and herbs have been used for many centuries to enhance the flavor and aroma of foods. Their importance has been recognized in preserving foods and for their medicinal value. Studies confirm that the growth of gram +ve and gram-ve food borne bacteria, yeast and moulds can be inhibited by Garlic (18) Cinnamon, Clove and other spices (18 , 19). The antimicrobial activity of plants is attributed to tannins, saponins, phenolic compounds, essential oils and flavanoids collectively known as

phytochemicals (15). Extracts of these herbs have proven potency against oral pathogens and are a genuine part of many of commercially available oral rinses and pastes.

Garlic is one of the greatest health tonics and has proven medicinal properties. It contains a substance called allicin which is equivalent to that of penicillin (1mg of allicin is equated to that of 15 IU of penicillin) (20). It also contains sulphur containing compounds like alliin, ajoene, diallylsulfide, dithiin, S-allylcysteine, enzymes, vitamin B, proteins and minerals. Also contains nonsulfur compound responsible for imparting antitumour, antioxidant and antimicrobial properties which can detoxify the body and has potent antioxidant activities. Allicin can destroy cell wall and cell membrane of root canal bacteria (21). Garlic inhibit the growth of oral pathogens such as *Streptococcus mutans* (21) and *P.gingivalis* and hence used in the management of dental infections such as periodontitis (21, 22). Studies report that a mouthwash containing 10% garlic in quarter Ringer solution produced a drastic reduction in oral bacteria (21).

Cinnamon which is a native plant of tropical islands is considered a herb and spice traditionally used by many ancient cultures. This plant has antimicrobial properties which has given it a huge prominence among its fellow herbs. Its bark as well as oil is both used to satisfy many antibacterial and antifungal requirements. Cinnamaldehyde is the organic compound that gives cinnamon its flavor and odour. It is especially effective against bacteria living at the back of the tongue reducing anaerobes by about 43%. It has also been found effective against *Streptococcus mutans* which is the causative organism of dental caries and also against *E.faecalis* (23).

Healing power of herbs still prevails in some cultures and one fourth of all modern medicine is derived from medicinal plants (Digoxin is derived from Foxgloves, Morphine and Codeine from Opium Poppies). Interest in medicinal herbs has grown during recent years. Studies using herbs as endodontic irrigants and medicaments are in progress and results are promising. The present study aims to establish antimicrobial efficacy of herbal extracts as endodontic irrigants.

Aims & Objectives

AIM

To evaluate the antimicrobial efficacy of Cinnamon and Garlic as endodontic irrigants against *E.faecalis* with comparison of 5.25% NaOCl and 2% CHX.

OBJECTIVES

1. To evaluate the antimicrobial efficacy of Cinnamon against *E.faecalis*.
2. To evaluate the antimicrobial efficacy of Garlic against *E.faecalis*.
3. To evaluate the antimicrobial efficacy of NaOCl against *E.faecalis*.
4. To evaluate the antimicrobial efficacy of CHX against *E.faecalis*.
5. To compare the antimicrobial efficacy of herbal extracts with NaOCl.
6. To compare the antimicrobial efficacy of herbal extracts with CHX.
7. To determine alternative herbal extract as endodontic irrigant against *E.faecalis*.

Review of Literature

R.R white et al (1997) (24) determined the residual antimicrobial activity of 2% CHX and 12% CHX after canal irrigation. After biomechanical preparation the canals were filled with sterile water and samples were collected using sterile paper points at 6,12,24,48 and 72 hrs. Antimicrobial activity was determined by determining the zone of inhibition in agar plates inoculated with *Streptococcus mutans*. Antimicrobial activity was present in all 2% CHX treated teeth throughout the 72hrs testing period and in most teeth, in relatively lower concentration for 6 to 24hrs after irrigation with 0.12% CHX.

L.B.Peters et al (2000) (25) developed a test model to quantify the penetration of bacteria into dentinal tubules. Model consists of two compartments separated by a bovine dentin specimen with pulpal side facing the incubated chamber of the test model. One compartment contained the test organisms such as *E.faecalis* and *A.israelii* and the other filled with sterile broth that was evaluated for growth of the test organism. The depth of bacterial penetration was measured with or without smear layer using both histological and quantitative recovering grinding technique after 6 wks of exposure. *E.faecalis* penetrated dentin significantly deeper than *A. israelii*. After removal of smear layer with EDTA, *E.faecalis* penetrated significantly deeper than in dentin penetrated with saline only or with a combination of saline and sodium hypochlorite.

Shyh – ming Tsao et al (2001) (26) investigated antibacterial activities of garlic oil and four diallyl sulphides against *Pseudomonas aeruginosa* and *Klebsiella pneumonia*. Antibiotic resistant profiles (Ceftazidime, Gentamicin, Imipenem, and Meropenem) were determined by using agar disc diffusion method. After 6hrs incubation the bactericidal effect of garlic oil against both organisms increased significantly with increase in concentration.

B.P.F.A Gomez et al (2001) (27) compared the antimicrobial activity of 0.5%, 1%, 2.5%, 4%, 2.5% NaOCl and 2 forms of chlorhexidine gluconate (gel and liquid) in three concentration (0.2%,1%,and 2%) in the elimination of *E. faecalis*. A broth dilution test using 24-well cell culture plates was performed and contact time taken for the irrigants to kill the bacterial cells was recorded. The tubes considered to have positive bacterial growth were confirmed by gram staining, colony morphology on BHI agar + blood and by use of biochemical identification kit. CHX liquid (0.2%) CHX gel (2%) produced negative cultures within 30s and 1minute respectively. When used at 0.2% concentration CHX gel destroyed bacterial cells after 2hrs of contact as opposed to 15minutes when used at 1% concentration. High concentration of NaOCl took less time to inhibit bacterial growth than lower concentration.

Bettina Basrani et al (2002) (28) evaluated the substantive antimicrobial activity of different medicaments such as 2% CHX gel, 0.2% CHX gel, 2% CHX solution + 2.5% CHX containing controlled release device for 7 days. After medication canals were inoculated with *E.faecalis* for 21 days. Dentin samples were collected with Gates-Glidden drills and bacterial growth were assessed with spectrophotometric analysis after 72 hrs of incubation. Mean optical densities were significantly lower for groups with 2% CHX when compared with those of controls. Canal dressing for 1 week with 2% CHX showed residual antimicrobial activity against *E.faecalis*.

J. C. Yamashita et al (2003) (29) evaluated in vitro the cleaning of root canal walls after irrigation with saline, 2% CHX, 2.5% NaOCl, 2.5% NaOCl + EDTA. The cleaning of the apical, middle and coronal thirds of the root canals was evaluated by scanning electron microscopy. The best cleaning was obtained using 2.5% NaOCl and EDTA followed by 2.5% NaOCl, whose cleaning was similar to CHX only in the

cervical third. Better cleaning was found in the cervical and middle third for all groups with the worst result in the apical third.

Onyeagba R. A .et al (2004) (30) studied antimicrobial effects of ethanolic extracts of garlic, ginger and lime against *Staphylococcus aureus*, *Bacillus* species, *E.coli* and *Salmonella* species. The entire test organism was susceptible to undiluted lime juice. The aqueous and ethanolic extracts of garlic and ginger did not inhibit any of these organisms. The highest zone of inhibition was observed with a combination of extracts on *Staphylococcus aureus*. *Salmonella* species were resistant to almost all the extracts except lime.

I. M. Saleh et al (2004) (31) investigated the ability of different endodontic sealers and Ca (OH)_2 to kill *E.faecalis* in experimentally infected dentinal tubules. After biomechanical preparation the canals were filled with GP and AH plus, Grossmans sealer, Ketac Endo, Apexit, Roekoseal Automix, Roekoseal Automix with an experimental primer or Ca (OH)_2 only. After 7 days the canals were reestablished with sterile Peeso reamers size 2. Dentin powder was removed using Peeso reamers size 5. Antimicrobial efficacy determined by reduction in CFU. AH Plus and Grossman sealer killed bacteria. Other endodontic sealers, as well as Ca (OH)_2 were less effective.

C. E. Radcliffe et al (2004) (32) evaluated the antimicrobial activity of varying concentrations of NaOCl such as 0.5%, 1%, 2.5%, 5.25% against endodontic microorganisms *A.israelii*, *A.naeslundii*, *Candida albicans*, *E.faecalis*. Contact time used was 0, 10, 20, 30, 60 and 120s. In case of *E.faecalis*, additional contact time such as 1.0, 2.0, 5.0, 10.0 and 30mts were tested. Pour plates were used to count low CFU and serial dilutions were used to find high CFU. All concentration of NaOCl lowered

the CFU below the limit of detection after 10s in case of *A.naesulundii* and *C.albicans*. *E.faecalis* proved significantly more resistant to NaOCl. But at higher concentrations less time was required though 5.25% NaOCl was not completely effective after 1minute.

Katharine. R. Carson et al (2005) (33) compared the antimicrobial activity of 6% and 3% NaOCl, 2% and 0.12% CHX, 0.01% and 0.005% doxycycline against four microorganisms such as *Prevotella intermedia*, *Streptococcus sanguis*, *Lactobacillus acidophilus*, *Peptostreptococcus micros* associated with primary endodontic infections. Agar diffusion test was used to determine the antimicrobial activities of the irrigants. MIC analysis was performed using macrodilution method. 0.01% and 0.005% doxycycline had significantly greater zones of inhibition than 6% NaOCl for *P.micros*, *P.intermedia*, *S.sanguis*. 6% NaOCl showed significantly greater zone of inhibition than 3% NaOCl for all endopathogens tested. 0.12% CHX had smaller zone of inhibition.

Fabio Roberto Dametto et al (2005) (34) evaluated the immediate and prolonged antimicrobial action of 2% CHX gel, 2% CHX solution, 5.25% NaOCl as endodontic irrigant against *E.faecalis* using dentinal tubule disinfection method. Controls were distilled water and natrosol gel. Samples were collected before mechanical preparation, after the preparation and 7 days after biomechanical preparation using sterile paper points. Antimicrobial efficacy determined by reduction in CFU. 2% CHX gel and liquid significantly reduced the CFU in the post treatment and 7 days after the treatment. 5.25% NaOCl reduced the CFU immediately but not able to keep the canal free of *E.faecalis* after 7 days.

Brian. D. Barnhart et al (2005) (35) evaluated the cytotoxicity of NaOCl, IKI, Betadine scrub, Ca(OH)₂, chlorine dioxide on cultured gingival fibroblast using the CYQuant assay. Human gingival fibroblasts were grown in Dulbeccos modified eagle medium containing 10% fetal bovine serum at 37°C and 5% CO₂. Cells were split plated for 24h to allow attachment. Irrigants were tested at different concentrations. IKI and Ca (OH)₂ were well tolerated by human gingival fibroblast.

Russell. S. Eddy et al (2005) (36) investigated the antibacterial efficacy of Chlorine dioxide (10% clidox -S), 13.8% Bioclenz, 5.25% Clorox, Saline against E.faecalis using dentinal tubule disinfection method. The dentin disk were incubated for 30 minutes and were then frozen, pulverized, serially diluted in phosphate buffered saline, and plated on BHI plates in triplicate. Antimicrobial efficacy determined by reduction in CFU. NaOCl was effective at eliminating E.faecalis than any of other irrigants. Bioclenz & Clidox-s were significantly better than saline, which was ineffective at eliminating E.faecalis.

Suree Nanasombat et al (2005) (37) investigated the anti bacterial activity of crude ethanolic extracts and essential oils of spices including cardamom, cinnamon, clove, corriander, cumin, garlic, ginger, holy basil, kaffir lime leaves and peels, lemongrass, mace, nutmeg, black and white pepper and turmeric against 20 sero types of Salmonella and 5 species of other Entero bacteria using disk diffusion method. The ethanolic extracts of clove, cumin and kaffir lime showed the broadest antibacterial activity while the extracts of cardamom, cinnamon, and kaffir lime leaves inhibited the growth of all strains except for S.typhimurium, S.london and Serratia marcescens. Oil of cloves and kaffir lime peels showed greater antibacterial activity against all strains tested.

Melissa L. Ruff et al (2006) (38) compared the antifungal efficacy of 6% NaOCl, 2%CHX, 17% EDTA and Biopure MTAD against *Candida albicans* inoculated root canals using dentinal tubule disinfection method. Aliquots from the experimental teeth were placed on Sabourauds dextrose agar plates and antifungal activity determined by reduction in CFU and significantly superior to biopure MTAD and 17% EDTA. MTAD was significantly superior to 17% EDTA.

M. S. Clegg et al (2006) (39) evaluated the effectiveness of 6%, 3%, 1%, 2% CHX, 1% NaOCl followed by Biopure MTAD, sterile phosphate buffered solution on apical dentin biofilms. Samples collected from patients were cultured on hemisections of root apices to generate a polymicrobial biofilm. Each biofilm was separately immersed in irrigants. SEM evaluation was done to determine the presence and morpholgy of bacteria. Dentin shavings were obtained using No 2 sterile carbide burs. Antimicrobial activity determined by reduction in CFU. 2% CHX was not capable of disrupting the biofilm. Viable bacteria could not be cultured from specimens exposed to 6% NaOCl, 2% CHX or 1% NaOCl followed by Biopure MTAD. 6% NaOCl was the only irrigant capable of both rendering bacteria and physically removing the biofilm.

Thomas .R. Dunavant et al (2006) (40) compared the efficacy of root canal irrigants against *E. faecalis* biofilm grown in a flow cell system. Tested irrigants were 6% NaOCl, 2% CHX, REDTA and Biopure MTAD. Antimicrobial efficacy was determined by reduction in CFU. No statistical difference was found between 1% and 6% NaOCl. A significant difference was found between 1% and 6% NaOCl and all other test agents such as 2% CHX, REDTA and Biopure MTAD.

Isabelle Portenier et al (2006) (41) investigated the antibacterial activity of MTAD and CHX towards two strains of *E. faecalis* and the inhibitory effect of dentin and bovine serum albumin (BSA) on the antibacterial activity. Survival of bacteria exposed to medicaments in the presence or absence of inhibitors was monitored in an in vitro model. Full concentration MTAD and 0.2% CHX killed both strains. Combining CHX with cetrimide further reduced the time required for killing. The presence of dentin or BSA caused a marked delay in killing by both medicaments.

V. B. Berber et al (2006) (42) evaluated the efficacy of 0.5%, 2.5% and 5.25% NaOCl as intracanal irrigant associated with hand and rotary instrumentation techniques against *E. faecalis* using dentinal tubule disinfection method. Sampling of the canal was done using three sterile paper points. Dentinal shaving were also collected by discing the tooth into three thirds and dentin chips collected using ISO 018, ISO 021, ISO 025, ISO 029 sterile diamond tipped conical burs. Antimicrobial efficacy determined by reduction in CFU. At all depths and thirds of the root canals and for all techniques used 5.25% NaOCl showed better antimicrobial efficacy than 2.5% NaOCl.

N.T. Sena et al (2006) (43) evaluated the antimicrobial activity of 2.5% and 5.25% NaOCl and 2% CHX gel and liquid against single species biofilm of *E. faecalis*, *Candida albicans*, *Prevotella intermedia*, *Porphyromonas gingivalis*, *Porphyromonas endodontalis* and *Fusobacterium nucleatum* generated on a cellulose nitrate membrane placed on agar medium. The microorganism was suspended using vortex and the inoculum was serially diluted (10-fold). Antimicrobial efficacy determined by reduction in CFU. Antimicrobial agents 5.25% NaOCl and 2% CHX killed the tested microorganism more rapidly. *P. intermedia*, *P. gingivalis* and *P. endodontalis* were

eliminated in 30s by all antimicrobial agents, with or without agitation in contrast with the facultative and aerobe strain.

Seung - Eunn yang et al (2006) (44) determined the effects of smear layer and CHX treatment on the adhesion of *E. faecalis* to bovine dentin using 40 dentin blocks. Blocks were immersed in a suspension of *E. faecalis* for 3hrs after treating the groups with saline (5mts), 17% EDTA (5mts), 2% CHX, (7 days), EDTA and CHX (7 days). The bacteria adhering to the dentin surface were counted by examination using a SEM. Significant amount of bacteria was retained on the samples treated with saline. Smallest amount of bacteria adhered to samples treated with 2% CHX for 7 days.

John. M. Williams et al (2006) (45) compared the detection of *E. faecalis* by Real Time quantitative PCR, Reverse transcription – PCR (RT-PCR), and cultivation during endodontic treatment. Intracanal samples were collected upon access, post-instrumentation /irrigation and post calcium hydroxide treatment from 15 primary and 14 refractory infections involving 29 single rooted teeth. The bacterium was up to 3 times more prevalent in refractory than primary infections at each collection step. Real time PCR detected significantly more *E. faecalis* positive teeth and samples than cultivation. Real time PCR and reverse transcription PCR are more sensitive methods than cultivation for detecting *E. faecalis* in endodontic infections.

Erika M Johnson et al (2006) (46) investigated the coaggregation interactions between *E. faecalis* clinical isolates & species such as *Peptostreptococcus anaerobius*, *Prevotella oralis*, *Fusobacterium nucleatum* & *Streptococcus anginosus* which were shown to induce apical periodontitis in monkeys. Coaggregation

interactions between *E. faecalis* & *F. nucleatum* play a major role in endodontic infections.

Bettina. R. Basrani et al (2007) (47) determined the minimum concentration of NaOCl required to form a precipitate with 2% CHX. X ray photon spectrometry and time of flight secondary ion mass spectrometry (TOF-SIMS) were used to qualify and quantify the precipitate. A color change and precipitate were induced in 2% CHX by 0.023% and 0.19% NaOCl respectively. Both XPS and TOF-SIMS showed the presence of para- chloroaniline in an amount directly related to concentration of NaOCl used.

Bradley. M. Newberry et al (2007) (48) compared the antimicrobial effect of 3% NaOCl , 17% EDTA and Biopure MTAD as final rinse on eight strains of *E. faecalis* using dentinal tubule disinfection method. No 4 sterile carbide round bur was used to prepare three sample of dentin shavings. MIC and MLC were calculated for each strains of *E. faecalis* using both tube dilution methods. MTAD inhibited most strains of *E. faecalis* growth when diluted 1:8192 times and killed most strains of *E. faecalis* when diluted 1:512 times.

Trisha. A. Krause et al (2007) (49) compared antimicrobial effect of Biopure MTAD , Doxycycline 100mg/ml, 10% Citric acid, 5.25% NaOCl in bovine tooth model against *E. faecalis*. Dentin shavings were collected using burs ISO 037 and 040. 10-fold dilution made & spiral plating done. Antimicrobial efficacy determined by reduction in CFU. Zone of inhibition was also determined for various dilutions of the samples by disc diffusion method. In bovine tooth model, doxycycline and NaOCl produced a significantly greater antimicrobial effect than other samples.

At deeper bur depth NaOCl remained significantly more antimicrobial. In all three dilutions doxycycline produced larger zone of inhibition.

Luciano Giardino et al (2007) (50) evaluated the antimicrobial efficacy of 5.25%NaOCl, Biopure MTAD ,Tetraclean against *E. faecalis* biofilm generated on cellulose nitrate membrane filters. Membrane filters were transferred into tubes containing antimicrobial solution test agent and incubated for 5, 30, 60mts at 20°C. Antimicrobial efficacy determined by reduction in CFU. 5.25% NaOCl was effective after 5mts when compared with other test group. Same effect was produced by Tetraclean only after 60mts. Tetraclean reduced 90% bacteria load after 5mts and >99.9% after 30mts. Bacterial load reduction using Biopure MTAD has not been significant after 5mts but much lower after 30mts than Tetraclean and 5.25% NaOCl.

Caio. C. R. Ferraz et al (2007) (51) compared the antimicrobial efficacy of 0.2%, 1% & 2%, CHX gel, 0.2%, 1% and 2% CHX solution, 0.5%, 1%, 2.5%, 4%, 5.25%, NaOCl against *S.Aureus* , *E.faecalis*, *S.Sanguis* , *S.Sobrinus*, *Actinomyces naeslundii*, *porphyromonas gingivalis*, *P. endodontalis*, *Prevotella intermedia* and *Prevotella denticola*. Antimicrobial efficacy determined by zone of inhibition. The largest inhibition zones were produced with 2% CHX gel being significantly different from the inhibition zones produced by all NaOCl concentration including 5.25%.

J. Craig Baumgartner et al (2007) (52) compared the antimicrobial efficacy of 1.3% NaOCl / Biopure MTDA to 5.25% NaOCl/ 15% EDTA against *E. faecalis* using dentinal tubule disinfection method. Samples were collected after instrumentation and irrigation using sterile paper points and were taken after additional canal enlargement. The investigation showed consistent disinfection of

infected root canals with 5.25% NaOCl/15% EDTA. The combination of 1.3% NaOCl/Biopure MTAD left nearly 50% of the canals contaminated with *E. faecalis*.

Mathew. J. Royal et al (2007) (53) compared the effectiveness of 5.25% NaOCl, MTAD and 2% CHX in the rapid disinfection of polycaprolactone –based root canal filling material (Resilon pellets). Resilon pellets and GP Pellets were contaminated with *E. faecalis* and disinfected with 5.25% NaOCl, MTAD and 2% CHX. Samples were placed in centrifuge tubes containing BHI agar and incubated. At 1, 3, 7 days broth were checked for turbidity and scored for growth. Gram staining was done to identify bacterial species. 5.25% NaOCl, MTAD and 2% CHX were effective in the rapid disinfection of Resilon and GP pellets and a 1minute immersion was sufficient for disinfection.

Monika Marending et al (2007) (54) evaluated the impact of different irrigation sequence of 2.5% NaOCl (exposure time 24 mts) and 17% EDTA (exposure time 3 mts) on the elastic modulus and flexure strength of standardized human root dentin bars prepared from extracted upper third molar. Specimens after exposure to irrigants were subjected to 3-point bending test using a universal testing machine. Modulus of elasticity and flexural strength were evaluated. 24hr exposure to NaOCl caused a significant drop in flexural strength compared with water or EDTA treated groups whereas the elastic modulus remain unaffected. Short exposure to EDTA did not affect the mechanical dentin parameters.

Jose. F. Siqueira et al (2007) (55) in this study assessed the bacterial reduction after chemomechanical preparation using 0.12% CHX as an irrigant and the additive antibacterial effect of intracanal dressing with a paste of Ca(OH)_2 in 0.12% CHX gel. Bacterial samples were taken from 13 patients with primary

intraradicular infections and chronic apical periodontitis. Bacterial samples were taken before treatment (S₁) after chemomechanical preparation using CHX (S₂) and after 7 day dressing with Ca(OH)₂ /CHX paste (S₃). Bacterial count and growth were determined by 16S Ribosomal RNA gene sequencing analysis. 0.12% CHX solution as an irrigant significantly reduced the number of intracanal bacteria but failed to render the canal free of cultivable bacteria in one half of the cases. Application of a 7 day intracanal dressing further increased significantly the number of cases yielding negative cultures.

William. J. Nudera et al (2007) (56) in this study determined the minimum inhibitory and bactericidal concentration of Triclosan and Triclosan with Gantrez against *Prevotella intermedia*, *Fusobacterium nucleatum*, *Actinomyces naeslundii*, *Porphyromonas gingivalis* and *E. faecalis*. The MIC of both test solutions was determined for each of the 5 microorganism using microtiter serial dilutions. MBC was also determined by streaking the samples on agar plates. Addition of Gantrez demonstrated bactericidal activity of Triclosan. Both Triclosan and Triclosan with Gantrez demonstrated bactericidal activity against the 5 specific endodontic pathogens.

Joshua. M. Davis et al (2007) (57) compared the antimicrobial effects of Dermacyn, Biopure MTAD, 2% CHX, 5.25% NaOCl against *E. faecalis*. Antimicrobial efficacy was determined by zone of inhibition using Agar disc diffusion method. Biopure MTAD showed a larger zone of microbial inhibition for both aerobic and anaerobic samples when compared with 2% CHX and 5.25% NaOCl. The zone of inhibition for 2% CHX and 5.25% NaOCl were not different from each other but resulted in larger zone of inhibition than Dermacyn and control (saline).

Sunita Bansod et al (2008) (58) evaluated the anti fungal activity of garlic oil, neem oil, jitra oil, lemongrass oil, dalchine oil((cinnamon), nilgri oil, clove oil, cardamom, mint oil, tulsi oil, ajwain, ashwagandha and zinger oil against aspergillus fumigatus and aspergillus niger. Maximum antimycotic activity was demonstrated by oils of Cymbopogon martini, Eucalyptus globulus and Cinnamomum zylenicum.

Zohreh Ahangari et al (2008) (59) compared the antimicrobial effects of 2.5% NaOCl, 2% CHX against E. faecalis contaminated root canals of human extracted teeth. SEM evaluation was done for 2 samples to confirm sufficient penetration of microorganism into dentinal tubules. After biomechanical preparation dentinal shaving were taken using H files and transported into test tubes containing BHI and incubated at 37⁰ for 96 hrs. Antimicrobial efficacy determined by reduction in CFU. In MTAD group, one sample showed bacterial growth. Bacterial growth was present in 5 samples in 2.5% NaOCl group and in 4 samples of CHX group.

P. Chivatxaranukul et al (2008) (60) investigated the dentinal tubule invasion and the prediliction of E. faecalis for dentinal tubule walls. The invasion of dentinal tubules in extracted human teeth by E. faecalis was measured ex vivo after 8 weeks of incubation. Extent and maximum depth of tubule invasion were assessed histologically and compared between groups. In adherence study vertically split root samples were prepared with longitudinally aligned dentinal tubules and fractured orthodentin. Samples were processed for analysis using SEM. Bacterial adhesion to tubule walls versus fractured orthodentin was calculated as number of cells per 100mm². The strain of E. faecalis showed moderate to heavy tubule invasion after 8wks. In adhesion studies more bacteria adhered to fracture orthodentin than to dentinal walls.

Md. Mahfuzul Hoque et al (2008) (19) compared the antimicrobial activity of ethanol, aqueous extracts, essential oils of cloves and cinnamon extract against food borne pathogens *Listeria monocytogenes*, *Staphylococcus aureus*, *E. coli*, *Salmonella enteritidis*, *Vibrio parahaemolyticus* and *Bacillus cereus* and 5 food spoilage bacteria. *Pseudomonas aeruginosa*, *P. putida*, *Alcaligenes faecalis*, and *Aeromonas hydrophila*. Ethanolic extract of clove was potentially active against *E. coli*, *L. monocytogenes*, *S. aureus*, *V. parahemolyticus*, *Pseudomonas*, *Aeromonas*, and *Alcaligenes faecalis*. The aqueous extract of clove was active against *S. aureus* strains and *V. parahaemolyticus*. The ethanolic extract of cinnamon was active only against *Staphylococcus aureus*. The essential oil of clove and cinnamon showed maximum inhibition for *A. hydrophila*.

Rachele Neglia (2008) (61) compared in vitro and ex vivo studies on the antibacterial activity of Tetraclean, a new generation endodontic irrigant, and NaOCl against *E. faecalis*. Bacterial activity assessed using membrane-filtration for Tetraclean and dilution- neutralization for NaOCl. Ex vivo study conducted using model of extracted and decoronated human teeth infected with *E. faecalis* and subsequently irrigated with either of the irrigant. Both irrigants display very similar bactericidal activity against *E. faecalis* in vitro. Ex vivo model show that only in teeth irrigated with Tetraclean, the bacterial burden gradually drop until no bacteria were detectable a few days post irrigation. In teeth irrigated with NaOCl the drop in bacterial burden was rapid but temporary and most of the teeth were colonized again by 48 hours post irrigation.

Sandeep Singh et al (2009) (62) compared the antimicrobial efficacy of Biopure MTAD and 2.5% NaOCl irrigation in infected root canals following single visit endodontic procedure. Single canal anterior and premolar teeth that had pulpal or periapical pathology were selected. Biomechanical preparation along with 2.5% NaOCl and 0.9% saline and Biopure MTAD was done for 5 minutes. Microbiological samples were taken using sterile paper points before and after biomechanical preparation. Antimicrobial efficacy determined by reduction in CFU. 2.5% NaOCl resulted in significantly greater number of anaerobic organisms when compared to Biopure MTAD.

S. Shobana et al (2009) (63) evaluated the antibacterial activity of two varieties of garlic against enteric pathogens such as *E. coli*, *Proteus mirabilis*, *Salmonella typhi*, *Shigella flexneri* and *Enterobacter aerogenes*. Ethanolic extracts of sativum was found to be highly effective against all the bacteria tested. *Enterobacter aeurogenosa* was not susceptible to aqueous extract of both garlic varieties.

Andrea Ardizzoni et al (2009) (64) evaluated the effectiveness of NaOCl, Tetraclean and MTAD against 54 *E. faecalis* clinical isolates by agar dilution assay. In addition using the dentinal tubule disinfection method compared the irrigation efficacy of both Tetraclean and MTAD. Antimicrobial efficacy determined by reduction in CFU. According to agar dilution assay 100% of the clinical strains inhibited by Tetraclean at 1:256 where as 4 clinical isolates still sensitive at 1:66536 dilution. In ex vivo model study using Tetraclean irrigation, an immediate and drastic drop to undetectable values of CFU was evident in 150 teeth. Teeth irrigated with MTAD, the bacterial counts dropped more gradually but steadily.

Ya Shen et al (2009) (65) in this study developed a biofilm model that closely mimicked in vivo biofilm and to determine its susceptibility to endodontic antimicrobial irrigants by con-focal laser scanning microscopy. Collagen coated hydroxyapatites and uncoated hydroxyapatite disk were inoculated with dispersed subgingival plaque for 3 wks. Biofilm were subjected to 1, 3 and 10 minute exposure to CHX plus and 2% CHX. The collagen coated hydroxyapatite biofilm was thicker than hydroxyapatite biofilm. Less bacteria were killed in C-HA biofilm than in HA model.

Rosina Khan et al (2009) (66) evaluated the antimicrobial activities of the crude ethanolic extracts of five plant *Acalia nilotica*, *Syzgium aromaticum* and *cinnamum zeylanicum*, *Eucalyptus globulus* against multidrug resistant strains of *E. coli*, *Klebsiella pneumoniae*, *Candida Albicans* and ATCC strains of *Streptococcus bovis*, *Pseudomonas aerogenosa*, *Salmonella typhimurium* and *Ecoli*. Extracts of *A.nilotica*, *C.zeylanicum* and *S aromaticum* showed the most potent activity against all the microorganism. *E.faecalis*, *S. aureus*, *S. bovins* and *S.mutans* were the most susceptible to all the plant extracts tested. *S. typhimurium*, *K. pneumoniae*, *E coli*, *p. aeruginosa* and *C.Albicans* were found to be sensitive to extracts of *A.nilotica*, *C..zeylanicum*, *S. aromaticum*.

Srinivasan Durairaj et al (2009) (20) evaluated the antibacterial activity and stability of aqueous extract of garlic against 17 multidrug resistant gram positive and gram negative bacterial isolates, including *Staphylococcus aureus*, *Salmonella typhi*, *Pseudomonas aeroginosa*, *E. coli* and *Proteus species* . Antibacterial activity determined by well diffusion method characterized by inhibition zones. The maximum zone of inhibition was observed in *Bacillus subtilis* and minimum for

Proteus species. Antibacterial efficacy maintained at room temperature was for maximum 7 days. At - 20⁰C the activity was maintained for 90 days.

Bishnu joshi et al (2009) (67) evaluated the antibacterial property of ethanolic extracts of different medicinal plants Ocimum sanctum, Cinnamomum zeylanicum, Xanthoxylum armatum and Origanum majorina against 10 medically important bacterial strains namely Bacillus subtilis , Bacillus cereus, Bacillus thuringiensis, Staphylococcus aureus, Pseudomonas species, Proteus species, Salmonella typhi, E. coli, Shigella dysentriae, Klebsiella pneumonia. Antibacterial activity was determined by agar well diffusion method. Among all the pathogens, all gram positive bacteria were inhibited by all 4 plant extract. Gram negative bacteria found to be resistant. Exceptionally Salmonella typhi showed zone of inhibition against extract of Ocimum sanctum.

Luciano Giardino et al (2009) (68) evaluated the antimicrobial effect of MTAD, Tetraclean, Cloreximid and 5.25% NaOCl on three common endodontic pathogen (E. faecalis, Porphyromonas gingivalis and Prevotella Intermedia). Antibacterial activity was determined by agar plate diffusion method. 5.25% NaOCl showed a high antimicrobial activity against anaerobic bacteria. MTAD and Tetraclean showed a high action against both, strictly anaerobic and facultative anaerobic bacteria. Chlorhexidine + cetrimide showed the lowest antibacterial activity against both facultative and strictly anaerobic bacteria.

Meenal. N. Gulve et al (2010) (16) compared the antimicrobial efficacy of ginger extract and 2% NaOCl against E. faecalis using agar diffusion method, antimicrobial efficacy determined by the zone of inhibition. Ginger extract showed significant inhibition compared with 2% NaOCl.

Ram kumar Pundir et al (2010) (18) evaluated the antimicrobial activity of ethanolic extracts of *Snzygium aromaticum* and *Allium sativum* against food associated bacteria (*Bacillus subtilis* , *B. megaterium* , *B. polymyxa*, *B. sphaericus*, *Staphylococcus aureus* , *E.coli*) and molds (*penicillium oxalicum*, *Aspergillus flavus* *A. luchuenis*, *Rhizopus stolonifer* , *Scopulariopsis species* and *Mulcor species*). Antimicrobial activity determined by agar well diffusion method. MIC& MBC was also determined. Garlic and clove showed antibacterial activity and zone of inhibition was in the range of 20 mm-31mm.

U. B. Owne Ureghe et al (2010) (69) studied the inhibitory activity of garlic and lime on seven bacteria isolated from 240 extracted carious teeth. Agar well diffusion method was used in the susceptibility testing and MIC was calculated. Considerable antibacterial activity of garlic and lime were noted against the seven bacteria (*Streptococcus mutans*, *Lactobacillus acidophilus*, *Nocardia asteroides*, *Pseudomonas aeruginosa*, *Actinomyces viscosus*, *Staphylococcus aureus* and *Veillonella alcaligenes*.

Vahid Zand et al (2010) (70) compared the efficacy of gel and solution forms of NaOCl in removal of smear layer from root canals. The canals of all teeth were prepared with rotary Race instruments and flushed with 2.5% NaOCl solution and in NaOCl gel group the canals were coated with gel & final rinse with 1ml of 17%EDTA for 2mts. The amount of smear layer was quantified according to the Torabinejad method using SEM. No significant difference between NaOCl solution and gel but there was significant difference between saline & NaOCl gel in the apical, coronal and middle third. Use of NaOCl gel can be effective as NaOCl solution along with EDTA in smear layer removal.

Carmen Maria Ferrer-Luque et al (2010) (71) evaluated the antimicrobial activity of Maleic acid and combinations of Cetrimide with chelating agents against E. Faecalis biofilm. E.faecalis biofilm grow in the MBFC high- through put device for 24hrs and exposed to irrigating solution. Maleic acid showed antimicrobial activity against E.faecalis biofilm both alone or in association with Cetrimide from 30s onwards and the combination of EDTA and citric acid with cetrimide eradicated biofilm after 1minute of contact.

Saurabh S .Chandran et al (2010) (72) evaluated the antifungal efficacy of 5.25% NaOCl, 2% CHX,17% EDTA with and without the inclusion of an antifungal agent (1% clotrimazole) against Candida albicans inoculated root canals using dentinal tubule disinfection method. Aliquots from the experimental teeth were cultured on Sabouraud agar and antimicrobial efficacy determined by reduction in CFU. 5.25% NaOCl exhibited superior antifungal efficacy compared with 2% CHX and17% EDTA. 5.25% NaOCl and 2% CHX with clotrimazole showed significantly greater antifungal properties than 17% EDTA with clotrimazole.

Andrea Cruz Camara et al (2010) (73) evaluated the antimicrobial activity of 0.2%, 1%, 2%CHX in root canals infected using dentinal tubule disinfection method against Candida albicans, Pseudomonas aeroginosa, E.faecalis and Staphylococcus aureus. Biomechanical preparation of the root canals was done using protaper universal system. Samples were taken using sterile paper points and antimicrobial efficacy determined by reduction in CFU. 0.2%CHX in combination with rotary instrumentation was ineffective against all test micro organism. The 1% CHX solution was ineffective against S.aureus and E.faecalis. The 2% CHX was not sufficient to inactivate E. faecalis.

Maria Teresa Arias-Moliz et al (2010) (74) assessed the antimicrobial efficacy of Cetrimide and CHX alone, in combination and alternating form in eradicating *E.faecalis* biofilm grown in the MBFC-high through put device for 24hrs. Contact time tested were 30s, 1 minute, 2 minute. Cetrimide eradicated *E.faecalis* biofilm at concentration of 0.5%, 0.0312% and 0.0078% at 30s, 1 minute & 2 minute contact times respectively. CHX did not eradicate the biofilm at any concentration or time. The associated use of Cetrimide and CHX provided better results than their applications as single agent and the alternating application was significantly more effective than the combined mode of application.

Luis E. Chavez de paz et al (2010) (75) in his study combined confocal microscopy, a mini flow cell system and image analysis to test in situ the effect of antimicrobials and alkali on biofilm of *E.faecalis*, *Lactobacillus paracaesi*, *Streptococcus anginosus*, *Streptococcus gordonii* isolated from root canals with persistent infections. Biofilm formed for 24hrs were exposed for 5mts to alkali, 2.5% CHX, 50mmol/L EDTA, 1% NaOCl. The biofilm were characterized by using fluorescent markers targeting all membrane integrity and metabolic activity. NaOCl affected the membrane integrity in all microorganisms and removed most biofilm cells. Exposure to EDTA affected the membrane integrity in all organisms but failed to remove more than a few cells in biofilms of *E.faecalis*, *L.paracasei*, *S.sanginosus*. CHX had a mild effect on the membrane integrity of *E.faecalis* and removed only 50% of its biofilm cells.

Bonnie Retamozo et al (2010) (76) investigated the concentration of NaOCl and the irrigation time required to disinfect dentin cylinders infected with *E.faecalis*. 1.3%, 2.5% and 5.25% NaOCl and contact time of 5, 10, 15, 20, 25, 30, 35 & 40mts were used. The visual turbidity model was used in the study to detect

remaining viable bacteria. Growth determination was done by formation of colonies in the agar plates. 1.3% or 2.5% NaOCl is ineffective in eliminating strains of *E. faecalis* at less than 40mts. 5.25% NaOCl was 100% effective at 40mts.

J. Prabhakar et al (2010) (77) evaluated the antimicrobial efficacy of Triphala, Green tea polyphenols, MTAD and 5% NaOCl against *E. faecalis* biofilm formed on tooth substrate. Zone of inhibition, MIC&MBC of test samples were also determined. At the end of third and sixth week the tooth with the biofilm formed were treated for 10 minutes with the test solution. Sample were taken by scraping the biofilm and inoculated on agar plates. Maximum inhibition was observed by 5% NaOCl (2mts) followed by MTAD (2mts) compared with Triphala and GTP. Three week biofilm showed complete inhibition of bacterial growth when treated with Triphala, MTAD and NaOCl, but samples treated with GTP and saline showed presence of bacterial growth. Six week biofilm showed growth when treated with Triphala, GTP and MTAD where as NaOCl showed complete inhibition.

F. G Pappen et al (2010) (78) investigated the antibacterial action of MTAD + 0.01% Cetrimide, MTAC-1 (Tween 80 replaced by 0.01% CTR in MTAD) MTAC-2 (Tween 80 replaced by 0.1% CTR) MTAD-D (MTAC without the tween 80 and no CTR) against planktonic cultures and mixed species in vitro biofilm model using direct exposure test. Tetraclean was more effective than MTAD against *E. faecalis* in planktonic culture and in mixed species in vitro biofilm. CTR improved the antimicrobial properties of the solution.

Sonja Stojicic et al (2010) (79) evaluated and compared the effects of concentration, temperature and agitation on the tissue dissolving ability of NaOCl. A hypochlorite product with added surface active agent was compared with

conventional hypochlorite Solution. Thus NaOCl solutions in concentrations of 1%, 2%, 4% and 5.8% were tested at room temperature 37°C and 45°C on bovine muscle tissue with and without agitation by ultrasonic and sonic energy and pipetting . Percentage of weight loss of tissue specimens were calculated before and after treatment and contact angle on dentin at concentration of 1% & 5.8%. Weight loss of the tissue increased with the concentration of NaOCl. The effect of agitation on tissue dissolution was greater than that of temperature; continuous agitation resulted in fastest tissue dissolution. Hypochlorite with added surface active agent had the lowest contact angle on dentin.

Isabela N. Rocas et al (2010) (80) studied the bacteria in endodontic treatment procedure by using a combined ribosomal RNA- based reverse transcriptase polymerase chain reaction and reverse capture checker board hybridization approach. Samples were taken from infected canals of teeth with apical periodontitis before treatment (S1) and after chemomechanical preparation with NaOCl & after medication with Ca (OH)₂ paste. Presence of bacteria was screened by DNA based single PCR assay. RNA extracts were subjected to RT-PCR and were surveyed for the presence of 28 targeted bacteria by checker board method. Bacteria were found in all (S1)) sample, detectable level of ribosomal RNA was seen in 60% of cases after chemomechanical preparation and 53% after intracanal medication.

Madhu Pujar et al (2011) (13) evaluated the antimicrobial efficacy of triphala, green tea poly phenols, 3% NaOCl against E.faecalis biofilm formed on tooth substrate. Teeth were prepared by sectioning them vertically and the concave tooth surface minimally grounded to achieve a flat surface to enable placement in tissue culture wells. Samples collected by scraping the biofilm from root canal portion and antimicrobial efficacy determined by reduction in CFU. 3% NaOCl showed

maximum antibacterial activity against 2 week *E.faecalis* biofilm. Triphala and GTPS showed significantly better antibacterial activity.

Moeen Mahmoud Al Weshah et al (2011) (81) investigated the effect of 2% CHX on the hardness of different levels of root dentin when used as root canal irrigant. 20 extracted teeth were endodontically treated and sectioned into 3 sections cervical, middle and apical and three sections were mounted on an acrylic resin disc shape mould. Microhardness was measured using a Wallace microhardness appliance at 5 locations at 1 mm distance from the root canal before and after irrigation with 2% CHX for an hour. No significant difference in the decrease of dentin hardness was noticed.

Zahed Mohammadi et al (2011) (82) evaluated the antibacterial substantivity of a new sodium hypochlorite based root canal irritant (HYPOCLEAN) Tetraclean, 5.25% NaOCl against *E.faecalis* infected root canals using dentinal tubule disinfection method. Dentin shavings were taken on 0, 7, 14, 21 and 28 days using sterile slow-speed round burs with increasing diameter of ISO sizes 025, 027, 029, 031 and 033 respectively. Antimicrobial efficacy determined by reduction in CFU. The substantivity of Tetraclean was significantly higher than both Hypoclean and NaOCl solutions and retained in the root canal dentin for at least 28 days. The modified NaOCl solution showed more effective antibacterial activity than 5.25% NaOCl at all experimental periods.

Poonam Shingare et al (2011) (83) evaluated the antimicrobial activity of 12.5% alcoholic extract of miswak, 11% alcoholic extract of propolis, 3% NaOCl and 4.09% saline as root canal irrigant in chronically exposed primary tooth. Pre and post irrigation samples collected using sterile paper points. Samples cultured on tryptose

soya agar and colonies counted using digital colony counter. Miswak could be a good natural substitute to NaOCl, while propolis showed results comparable to that of saline.

Ashish Saraf et al (2011) (84) evaluated the antibacterial activity of two different Cinnamon species *C.zeylanicum* (commercial variety) and *C.flexuosus* against human pathogenic bacteria *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Bacillus subtilis*. Maximum inhibition was shown by *C.zeylanicum* against gram positive bacteria *S.aureus*. *C. flexuosus* showed inhibition but was inactive against *Klebsiella pneumoniae*. *Bacillus subtilis* was least inhibited.

Zohreh Dalisrani et al (2011) (85) compared the antimicrobial effect of chamoline, garlic, clove, cinnamon, sage, thyme against *Candida albicans*. Non growth halo in disks containing chamoline, garlic, clove, cinnamon, sage and thyme was observed. Cinnamon, garlic, chamoline, sage, clove, thyme had inhibitory effects on *Candida albicans*.

Mehrdad Lotfi et al (2011) (86) compared the antimicrobial efficacy of nanosilver (NS), NaOCl and CHX against *E. faecalis*. The minimum inhibitory concentration was determined by serial dilution method. Zone of inhibition was determined by agar disc diffusion method. The concentration used were 5.25% NaOCl, 0.33% NaOCl, 25µg/ml of NS (0.0025%) 50 µg /ml (0.005%) of NS, 4000 µg/ml (0.4%) of NS and 2% CHX. Vancomycin bacterial susceptibility papers were used as positive and sterile saline as negative control. Zone of inhibition for 2% CHX were significantly larger than other groups. 0.0125% NS showed inhibition effect against *E.faecalis* in MIC test in the first 6 hrs, whereas Naocl could exert the same

effect at a concentration of 0.082%. Concentration of NaOCl should increase approximately 70 fold to demonstrate the same antibacterial effect as NS.

Melahat Gorduysus et al (2011) (87) compared the antimicrobial effects of various endodontic irrigants 2% NaOCl, 2% CHX, 2.4% Iodine pottasium iodide, 17% EDTA and bioactive glass, against four selected microorganisms *E. faecalis*, *E. coli*, *S. aureus*, *S.pyogenes*. MIC and MBC were determined. 17% EDTA was most effective against *S.pyogenes*, *S.aureus*, *E. faecalis* in 24 hrs. NaOCl displayed activity equivalent to that of IKI on *E.coli*, *S. aureus* and *E. faecalis*.

M N Shahani et al (2011) (88) compared the antimicrobial substantivity of 2% CHX, 1% Povidone iodine, 2.5% hydrogen peroxide followed by 2% NaOCl alone as irrigants in instrumented root canals up to 72 hrs. After biomechanical preparation of the canals endodontic paper point were placed for 2mts and removed and stored in cryogenic vials at refrigerator temperature. Procedure was repeated after 12, 24, 48 & 72 hr. Antimicrobial activity determined by zone of inhibition. 2% CHX showed antimicrobial substantivity lasting for 72 hr followed by 1% Povidone iodine and 2% NaOCl.

Ponmurgan Karuppiah et al (2012) (89) evaluated the antibacterial effect of allium sativm, clove and zingiber officinale against multidrug resistant clinical pathogens causing nosocomial infection. All bacterial isolate were susceptible to crude extracts of both plant extracts. Except enterobacter species and klebsiella species, all other isolates were susceptible. The highest zone was observed with garlic against *Pseudomonas aeruginosa*.

D. Mortenson et al (2012) (90) determined the effect of using an alternative irrigant between NaOCl and CHX to prevent the formation of parachloroaniline

within the root canal system. After biomechanical preparation the samples were irrigated with saline followed by 2%CHX, 50% citric acid followed by 2% CHX, 14% EDTA followed by 2%CHX. The chemical identity and quantification of the PCA in the formed precipitate was determined using gas chromatography/ mass spectrometry. All experimental group contained PCA. Citric acid used as the intermittent irrigant had the least amount of PCA formation in the canal system.

S. Stojicic et al (2012) (91) evaluated the antibacterial efficacy and smear layer removal of a novel root canal irrigant QMix against *E.faecalis* and mixed plaque bacteria in planktonic phase and biofilm . *E.faecalis* and mixed plaque bacteria were exposed to QMix, 2% CHX, MTAD, 1% NaOCl for 5s, 30s and 3mts. Samples serially diluted and cultured aerobically and anaerobically to measure killing of bacteria. The amount of killed bacteria in biofilm on dentin disc was analyzed by confocal laser scanning microscopy. The effectiveness of smear layer removed was evaluated by SEM. QMIX and 1% NaOCl killed all planktonic *E. faecalis* and plaque bacteria in 5s while 2% CHX and MTAD were unable to kill all plaque bacteria in 30s. QMIX and 2% NaOCl killed up to 12 times more biofilm bacteria than 1% NaOCl and MTAD. QMiX removed smear layer equally well as EDTA.

Materials & Methods

MATERIALS

K- FILES	- MAILLEFER, DENTSPLY
5.25% NaOCl	- NICE CHEMICALS Pvt, Limited.
2%CHX	- VIJAY DENTOCARE
17%EDTA	- VISTA DENTAL PRODUCTS
Normal saline	- BAXTER Pvt. Limited
Broaches	- MILTEX COMPANY
Hedstrom files	- MAILLEFER, DENTSPLY
Gates gliddden drills	- MANI
Diagnostic sensitivity agar	- HI - MEDIA, MUMBAI
ATCC 29212 strains of E.faecalis	- HI- MEDIA, MUMBAI
Streptococcus selection broth	- HI -MEDIA, MUMBAI
Sheep Blood Agar	- HI - MEDIA, MUMBAI
McConkeys Agar	- HI- MEDIA, MUMBAI
Petridishes	- BOROSIL COMPANY
Micropipettes	- BOROSIL COMPANY
Test tubes	- BOROSIL COMPANY
Freshly prepared aqueous extract of Cinnamon and garlic	-
PANKAJAKASTHURI RESEARCH CENTRE, TRIVANDRUM.	

EQUIPMENTS USED

Airotor hand piece	- NSK PANA-AIR
Micromotor handpiece	- DTMUSA, NSK
Autoclave	- All AMERICAN 41 QUART ELECTRIC STERILIZER MODEL# 75X
Steam distillation unit	- CORNING COMPANY
Rotator	- REMI MAKE COMPANY
Water bath	- BESTON MAKE COMPANY
Incubator	-BESTON MAKE COMPANY
Laminar flow	- ROTEK COMPANY

SELECTION OF TEETH AND CANAL PREPARATION

Fifty human mandibular premolar with roots of similar form were selected for the study of which specimens selected were from teeth after orthodontic extraction. The teeth selected had single canal with straight roots measuring approximately 21mm. The teeth with caries, crack and restorations were excluded from the study. All the external debris were removed from the tooth surface with an ultrasonic scaler and teeth were placed in 0.5% NaOCl for 24hrs for surface disinfection and were stored in normal saline until use.

In the first step of the study, the anatomical crown of all the teeth were cut away at Cemento enamel junction(CEJ) perpendicular to the long axis of the teeth using a water cooled diamond wheel bur on an airturbine hand piece at 300000 rpm. The remaining roots measured 12-18 mm. The exploration of the radicular canal was accomplished with no 10 and no 15K file to make sure that the roots had only one canal and it was patent. The preparation of the entrance of radicular canal was done with rotary instrument gates glidden drills of sizes 2 and 3. An appropriate sized barbed broach was used to extirpate the pulp.No15 size K file was inserted into the canal so that the tip of the file was visible at the apical foramen. Then the working length was determined one mm short of the file penetration into the canal. K files of sizes 15-50 were used to biomechanically prepare the root canal. The canals were recapitulated and irrigated with 5.25% NaOCl, 2% CHX, 17% EDTA and final rinse was with normal saline. Subsequent to the canal preparation the apical foramen of all the specimens were sealed with cyanoacrylate glue to prevent bacterial microleakage.

Specimens were placed in steel containers containing BHI broth and subjected to autoclave at 121°C at 15 PSI for 20 minutes for sterilization. Subsequent to sterilization all the specimens were transported and manipulated under laminar flow using sterile instruments and equipments.

PREPARATION OF E.FAECALIS SUSPENSION

In order to get a controlled and standard suspension of the organism the following procedure was adopted:

From a stock culture of ATCC 29212 E.faecalis strain, subculture was made onto a plate of Diagnostic Sensitivity Test Agar. From this a typical colony was sub-cultured into 50 ml of Streptococcus Selection Broth contained in a 100 ml conical

flask. This was incubated at 37° c for 24 hours. Enumeration of live bacteria (CFU) was carried out as follows:

1 ml of the broth culture was added to 9 ml of sterile normal saline and mixed well to get a dilution of 1 in 10 (10^{-1}). The broth culture in the conical flask was kept in the refrigerator at 8° C. From the 10^{-1} dilution 1 ml was added to 9 ml of sterile normal saline to get a dilution of 1 in 100 (10^{-2}). This serial dilution was continued until a dilution of 10^{-6} . 1 ml from each dilution was pipetted into a sterile petri dish in duplicate. 20 ml of Diagnostic Sensitivity Agar melted, cooled and was added to each plate, mixed well and allowed to solidify. The plates were incubated at 37°C for 24 hours. The plates were examined for growth. The plates in which the number of colonies is between 30 and 300 were selected. The average number of colonies of duplicate plates was calculated. This number was multiplied by the dilution factor to get the number of colony forming units in 1 ml of the original broth culture. The average number of CFU of the dilution 10^{-5} was 38. The concentration of the original broth culture was computed to 3.8×10^5 CFU per ml. For injecting into the tooth a suspension of bacteria containing 10^5 CFU per ml was used. For preparing this 10 ml, original broth culture was diluted to 38 ml with sterile normal saline.

AQUEOUS EXTRACTION PROCEDURE

100g of air dried, coarsely powdered plant material was taken in a 1 litre round bottom flask and 400ml of water added to the sample. The flask was fitted with a water condenser and the mixture was refluxed for 2 hours, cooled and filtered. The filtrate was evaporated over a water bath to 100 ml.

TOOTH INOCULATION WITH E.FAECALIS

The root canals were inoculated with E.Faecalis suspension using sterile 1ml tuberculin syringes and specimens were separately placed in steel containers containing 2ml of broth. The steel containers containing the specimens were kept in incubators at 37°C for 21 days. After 21 days all the specimens were retrieved and each specimen was transferred into test tubes containing 3ml of saline and was shaken three times for 30 seconds each time on a rotator to remove the excess culture medium. In addition large amount of bacteria present on the surface of the specimen were removed during rinsing and irrigation. One sample was subjected for SEM evaluation to confirm the penetration of microorganism into the dentinal tubule.

The contaminated samples were divided into five groups, each containing ten teeth. Test irrigating solutions were used as follows.

Group 1 - Normal Saline

Group 2 - 5.25% NaOCl

Group 3 - 2% CHX

Group 4 - Cinnamon

Group 5 - Garlic

Groups were irrigated with respective irrigating solutions using 2 ml syringes and were immersed in test tubes containing 2ml of the solution for 5mts. Subsequent to the removal of specimens from the test tubes each specimen was transferred into test tube containing 3ml of saline and shaken in a rotator for 3 times for 30 second

each. Dentinal shavings were collected using no 40 H file in an aseptic condition. Shavings were transferred into test tubes containing 10 ml sterile normal saline (10^{-1}). One ml from the above was diluted to 10 ml with sterile normal saline (10^{-2}). One ml from the above was diluted to 10 ml with sterile normal saline (10^{-3}). One ml from the above was diluted to 10 ml with sterile normal saline (10^{-4}). One ml from the above was diluted to 10 ml with sterile normal saline (10^{-5}). One ml from the above was diluted to 10 ml with sterile normal saline (10^{-6}).

Three dilutions from the above viz., 10^{-1} , 10^{-3} and 10^{-6} were used for the count. From this one ml from each dilution was pipetted on to a sterile 100 mm diameter in duplicate. To each of these plates 15 ml of agar medium, melted, cooled and was added mixed well and allowed to solidify. These plates were incubated for two days at 37°C . After incubation the number of colonies was counted in suitable plates. The number of the colonies multiplied by the dilution factor gives the total number of CFU in the scrapings per tooth.

TO CHECK WHETHER CONTAMINATION BY ANY OTHER BACTERIA HAS NOT TAKEN PLACE

In order to ascertain that contamination by any other bacteria has not taken place during the procedures the following test were carried out:

From the initial 1 in 10 dilution of the scrapings from each of the tooth sub cultures were made using a bacteriological loop onto (1) a plate of sheep blood agar and (2) a plate of McConkey's agar plate. These plates were incubated at 37°C for 24 hours. After incubation the growth in the plates were examined for their colonial morphology and by Gram's staining. Growth in all the plates was of typical

morphology of *E. faecalis*. Only one type of colonies was found on each plate. By Gram's staining typical gram positive cocci in chains were observed.

Results & Observations

Antimicrobial activity was determined by evaluating the CFU in three dilutions 10^{-1} , 10^{-3} , 10^{-6} . The no of viable colonies and percentage of inhibitory effect was recorded in different dilutions, tabulated and are shown in tables 1, 2, 3, 5, 6, 7.

STATISTICAL ANALYSIS OF THE RESULTS

The statistical analysis was performed using SPSS software (Statistical Package for Social Sciences) version 16.0. The data was interpreted at a confidence interval of 95%. Analysis of variance (ANOVA) was used to compare the antimicrobial activity of herbal extracts with comparison of standards. Post Hoc test followed by Scheffe Test was performed for multiple comparisons of the specimens.

(P value < 0.05 considered as significant difference).

TABLE 1 shows mean values of number of viable colonies of bacteria in each group at 10^{-1} dilution.

The statistical test shows significant difference in the multiple comparison between the groups but there is no significant difference between group 2 compared with group 3 at 0.05 (95% confidence interval)

TABLE 2 shows mean values of number of viable colonies of bacteria in each group at 10^{-3} dilution.

The statistical test shows significant difference between group1 and other groups, but there is no significant difference between groups 2, 3, 4, 5 in the multiple comparisons.

TABLE 3 shows mean values of number of viable colonies of bacteria in each group at 10^{-6} dilution.

The statistical test shows significant difference between group1 and other groups, but there is no significant difference between groups 2, 3, 4, 5 in the multiple comparison groups, but no significance between group 2 and group 3.

TABLE 4 shows multiple comparison of mean number of viable colonies of bacteria in the different groups.

The statistical test shows significant difference in the multiple comparison between the groups but there is no significant difference between group 2 compared with group 3 at 0.05 (95% confidence interval)

TABLE 5 shows mean values of number and percentage of samples with growth of bacteria in each group at 10^{-1} dilution.

The statistical test shows significant value comparing group 1 with other groups, but no significance between group 2 and group 3.

TABLE 6 shows mean values of number and percentage of samples with growth of bacteria in each group at 10^{-3} dilution.

The statistical test shows significant value comparing group 1 with other groups, but no significance between group 2 and group 3.

TABLE 7 shows mean values of number and percentage of samples with growth of bacteria in each group at 10^{-6} dilution

The statistical test shows significant value comparing group 1 with other groups, but no significance between group 2 and group 3.

TABLE 8 shows multiple comparisons of mean values of number and percentage of samples with growth of bacteria in between groups

The statistical test shows significant difference when comparing group 1 with other groups, group 2 with other groups, group 3 with other groups, group 4 with other groups, group 5 with other groups.

GRAPH 1 The statistical test shows significant value comparing group 1 with other groups, but no significance between group 2 and group 3.

GRAPH 2 The statistical test shows significant difference in the multiple comparison between the groups but there is no significant difference between group 2 compared with group 3 at 0.05 (95% confidence interval)

GRAPH 3 The statistical test shows significant difference when comparing group 1 with other groups, group 2 with other groups, group 3 with other groups, group 4 with other groups, group 5 with other groups.

Interpretation of Results

Standard irrigants showed complete inhibition in 10^{-1} , 10^{-3} , 10^{-6} dilutions. Test irrigants showed inhibition in all the dilutions. Saline showed growth in 10^{-1} , 10^{-3} , 10^{-6} dilutions. CHX and NaOCl showed complete inhibition in all the samples tested. Cinnamon and Garlic showed inhibition in 1 and 2 samples respectively in 10^{-1} dilutions. Among the herbal extracts garlic was more effective than Cinnamon.

Discussion

Long term success in endodontic treatment depends on complete debridement and disinfection of the pulp space. Despite a thorough mechanical preparation pulp remnants, debris, bacteria may be present in the irregularities of root canal system (92). Hence it is highly desirable that instrumentation should be accompanied by irrigation and intracanal medicaments for the elimination of microorganisms (4). In this present study a modification of Haapasalo and Orstavik model was used for assessing the antimicrobial efficacy of endodontic irrigants in dentinal tubule disinfection (5). The model was further modified by using extracted human teeth rather than bovine teeth. This modification was appropriate because of the marked difference in diameter between the canals of bovine and human teeth (28). Quantitative estimation of CFU in the dentinal tubules after disinfection was included in the present study which was a modification of Haapasalo and Orstavik model (93).

Persistent endodontic infections are mainly due to retention and recolonization of microorganisms in the dentinal tubule (7). *E. faecalis* was selected as the test organism because it is a facultative organism that is non-fastidious, easy to grow and efficiently and rapidly colonizes the tubules (94). It has been used extensively in endodontic research because it has been detected in 63% of post treatment diseases (95) and due to the high level of resistance to a wide range of antimicrobial agents. *E. faecalis* has been used as the test organism to determine the efficacy of endodontic irrigants in the previous in vitro studies. ATCC 29212 strains of *E. faecalis* which has been the standard strain used in the previous studies was selected for the present study (96).

Orstavik and Haapasalo proved that *E. faecalis* penetrated entire width of circumpulpal dentin just within 2 days of incubation. In their study bacteria was allowed to penetrate from both pulpal and periodontal side (94). A longer time is

needed for the organism to penetrate deeper into root dentin and there was delayed bacterial penetration when root cementum is left intact (97). In this study the cementum was left intact to simulate clinical conditions (36). The minimum instrumentation size required for the penetration of irrigants in the apical third is 30 (98). The canals were enlarged up to 50 K file in the present study. *E.faecalis* can penetrate dentinal tubules to a depth of 300-400 μm within 3 weeks. Prolonged incubation period increased the number of infected dentinal tubules but depth of penetration of bacteria increases slowly with time (5). Hence in this study the teeth were inoculated with the organism and incubated for 21 days.

Ingrowth or progress of bacteria into the dentinal tubules could be delayed or prevented by the presence of a smear layer. Etching of dentin before exposure results in deeper penetration (99). 17% EDTA was used in this study for removing the smear layer in the experimental specimens before autoclaving and inoculation. Another important factor for the survival of bacteria is the availability of a nutrient source (100). The teeth were immersed in the streptococcus selection broth and the broth was replaced on alternate days during the 21 day incubation period. Subsequent change of the broth allowed the microorganism to rearrange in biofilms which is a structure known to confer resistance of microbial cells to different antimicrobial agents (40). Another reason for the replacement of the broth was to avoid medium saturation (28). Most of the previous studies evaluated the presence of bacteria using SEM or light microscopy (101). In the present study SEM evaluation was done to confirm the presence of bacteria.

In most of the in vitro studies sample preparation was done using burs of different sizes, Gates Glidden drills, Hedstrom files and paper points. In this study sample preparation was done using 40 size H files which was similar to the sample collection

done in a previous study in 2007 (102). The methodology adopted for enumeration of CFU (Pour plate method) was in resemblance with the study conducted by Lynn et al (93).

For many years irrigants have been used as an adjunct to enhance the antimicrobial effect of cleaning and shaping in endodontics. Sodium Hypochlorite is a widely used irrigant as it covers most of the ideal properties of an endodontic irrigant especially the tissue dissolving property. Studies prove that 5.25% NaOCl has been widely used for many years and it prevails as the golden standard today (11, 12). Interaction of NaOCl with the tissue fluids, blood, dentin and other organic debris can reduce the effectiveness. Chemomechanical preparation is a short term procedure and NaOCl remains in the canal for only a few minutes. So the antimicrobial effectiveness of NaOCl within the root canal is a function of concentration and contact time (103). Studies regarding the concentration of NaOCl ranging from 0.5% to 5.25% has been conducted in the past. Concerns regarding cytotoxicity dependent concentration of NaOCl have led Bystrom and Sundqvist (1983) recommend 0.5% NaOCl as the ideal concentration with antimicrobial action (4). But this is in contradiction to the study conducted by Vianna et al 2004 which conclude that reduction in the concentration of 5.25% NaOCl decreased antimicrobial effectiveness against the anaerobes tested(104). In the present study a contact time of 5 minutes was taken as the standard time for all the irrigants. This could be explained on the basis of maximum antibacterial action exhibited by Cinnamon and Garlic in a 5 minute contact time during the pilot study. Results from previous studies have shown that 5.25% NaOCl can eliminate *E.faecalis* in a short exposure time of less than 30 seconds (27), 30 seconds (43) and 2 minute (32) which contradicts the findings of the present study. The difference in contact time may be attributed to the following factors.

1. In the previous study there is direct contact of microorganism with the antimicrobial agent, bacterial suspension were mixed with the antimicrobial agent where as in the present study the dentinal tubules are inoculated with the organism to simulate the clinical condition.
2. The inhibitory effect of dentin on the bactericidal effect of the irrigant has been documented (105).

An important limitation of many studies when evaluating the endodontic microbiota refers to sample preparation. In comparison to the study conducted by Berber et al 5.25% NaOCl eliminated ATCC 29212 strains of *E.faecalis* in a 10 minute contact time (42). The methodology adopted in the latter study is similar to the present study except for the sample preparation. Burs of different sizes were used to procure dentinal shavings unlike H files used in the present study. Bacterial inhibition at greater depths was assessed using burs of different sizes whereas H files were indicative of bacterial inhibition to a limited depth. Increasing the concentration is undesirable because it is an irritant to periapical tissue (106). Other undesirable effects of NaOCl are its unpleasant taste, high toxicity, corrosive to instruments, reduces the elastic modulus and flexural strength of dentin (13).

2% Chlorhexidine digluconate is another standard irrigant tested in this study. Chlorhexidine is a cationic bisguanide that act by adsorbing onto the cell wall of the microorganism causing the leakage of intracellular components. At low concentration CHX is bacteriostatic due to leakage of small molecular weight substances such as potassium and phosphorous. At high concentration it has bactericidal effects due to precipitation or coagulation of cytoplasm caused by protein crosslinking (107). Studies prove that 2% CHX has good antimicrobial action (108). In the present study 2% CHX produced complete inhibition of *E.faecalis* in 5 minute contact time which is

evidenced by the studies conducted by Oncag et al 2001 proving that 2% CHX was effective against *E.faecalis* in a 5 minute contact time (109). Another important property of 2% CHX is substantivity (110). A study proved that 2% CHX application after 10 minutes of application produced antibacterial effects for up to 12 weeks (111). Though it has broad antimicrobial activity, it lacks tissue dissolving property. Allergic reactions such as contact dermatitis, desquamative gingivitis, discolouration of teeth and tongue and dysgeusia have also been reported (12). Studies have evaluated the genotoxicity and have proved that 2% CHX is biocompatible.

Even after thorough chemomechanical preparation bacteria can still be recovered from canals (6). To combat the emerging antimicrobial resistance and considering the undesirable side effects of synthetic drugs, herbal alternatives might prove to be advantageous. Approximately 60% to 80% of the world population relies on traditional medicines for the treatment of common illness (112). Medicinal plants are known to produce certain bioactive molecules which react with other organisms in the environment inhibiting bacterial or fungal growth (113). Contrary to synthetic drugs antimicrobials of plant origin are not associated with many side effects and have an enormous therapeutic potential to heal many infectious diseases (112). An important characteristic of plant extract and their components is their hydrophobicity which enabled them to partition the lipids of bacterial cell membrane and mitochondria disturbing the cell structures and rendering them more permeable. Extensive leakage from the bacterial cells or the exit of critical molecules and ions will lead to death (67).

Serial dilutions are used when initial concentration of bacteria is too high to perform a plate count. 30-300 CFU are ideally required to use this method. In this study normal saline was used as diluents to provide ions for the survival of bacteria

and not the nutrients (114). Test irrigants Cinnamon and Garlic showed reduction in CFU in all the samples tested at 10^{-1} dilution. Complete inhibition was observed in 2 samples of garlic and 1 sample of Cinnamon. These results are indicative of potent antimicrobial action by the test irrigants. Saline showed innumerable colonies in the same dilution. Comparing the antimicrobial activity of herbal extracts Garlic had fewer colonies than Cinnamon and hence more antibacterial.

Garlic (*Allium Sativum*) is a bulbous perennial medicinal plant which belongs to the family Liliaceae. The antimicrobial activity is attributed to thiosulphinates. Studies reported that if extract is free from thiosulphinates the antimicrobial activity will be lost (113). Garlic can be used on microorganism that has particularly developed resistance to antibiotics. This is concluded in a previous study (2001) which explained that garlic oil and 4 diallyl sulphides showed in vitro activity against antibiotic resistant *Pseudomonas aeruginosa* and *Staphylococcus aureus* (115). Multidrug resistant *E.faecalis* can be made vulnerable by the medicinal properties of garlic. Studies proved that extracts of garlic are bactericidal and are effective against *E.coli*, *S.aureus*, *B.cereus*, *Salmonella*, *listeria*, *proteus* and *Streptococcal* species (116). Garlic contains many sulphur containing compounds and one of which is the odourless molecule alliin. When the plant is cut or damaged alliin comes into contact with the enzyme alliinase, which converts alliin into allicin. The later is responsible for the typical smell and medicinal properties of garlic. In microorganisms allicin interferes with lipid synthesis and RNA production. The target enzyme with which allicin interacts has been identified as acetyl-CoA Synthetase. Studies have proved that allicin has antifungal and anti thrombotic effects (116).

Cinnamon (*Cinnamomum zeylanicum*) is primarily used for its aromatic bark as a spice. The antibacterial activity of cinnamon may be attributed due to the presence

of cinnamaldehyde compound which inhibits the amino acid decarboxylation activity in the cell which leads to energy deprivation and microbial cell death (113). Antibacterial activity of two essential oils *C.zeylanicum* and *C.flexuosus* against gram negative and gram positive organism has also been proved (84).

Test irrigants used in this study was not adequate to classify herbal extracts as antimicrobial agents against *E.faecalis*. The active compounds imparting the antimicrobial effect of Cinnamon and Garlic would have been assessed if a chromatography was performed. To recommend herbal extracts as endodontic irrigants, it has to satisfy all the ideal properties of an irrigant. But in the present study only the antimicrobial property has been studied. With a deeper knowledge of other properties of herbal extracts and if found to satisfy the properties of an ideal irrigant it could open doors to a new era in endodontics.

Summary & Conclusion

Complete debridement and disinfection of the pulpal space is considered to be essential for predictable long term success in endodontic treatment. Eliminating microorganism from root canal system is possible only by a thorough chemomechanical preparation. However complete sterilization of pulp space is not always achieved due to extremely complex anatomy. *Enterococcus faecalis* is the most common species isolated in root filled teeth with apical periodontitis. The constant increase in antibiotic resistant strains and side effects caused by synthetic drugs has prompted researchers to look for herbal alternatives. The purpose of this study was to evaluate the antimicrobial efficacy of Cinnamon and Garlic as endodontic irrigants against 5.25% NaOCl and 2% CHX. Extracted human teeth were biomechanically prepared, autoclaved and inoculated with *E.faecalis* and incubated for 21 days. After 21 days the teeth were randomly divided into five groups and treated with respective irrigants for 5 minutes. Dentinal shavings were collected using H files. Antimicrobial efficacy was indicated by reduction in CFU (Pour plate method). 5.25% NaOCl and 2% CHX showed complete inhibition in all the samples tested. Garlic and Cinnamon showed reduction in CFU in all the samples tested and there was complete inhibition in 2 samples of Garlic and 1 sample of Cinnamon. The results are indicative of potent antimicrobial action. The use of herbal alternatives as a root canal irrigant might prove to be advantageous considering the several undesirable characteristics of standard irrigants and other frequently used antimicrobials. Further research is warranted to conclusively recommend herbal solutions as a root canal irrigant.

Standard irrigants 5.25% NaOCl and 2% CHX showed complete inhibition and remain as the standard irrigants. With ever increasing resistance to synthetic drugs and typical features of *E.faecalis*, herbal extracts can be an alternative option provided all the ideal properties of an irrigant are satisfied.

Table-1: Mean values of number of viable colonies of bacteria in each group at 10^{-1} dilution

Group	Number of viable colonies (MEAN\pmSD)
Group-I	1057.72 \pm 1.24
Group-II	0.00 \pm 0.00*
Group-III	0.00 \pm 0.00*
Group-IV	22.00 \pm 1.41* ^{#,†}
Group-V	15.00 \pm 1.00* ^{#,†}

Table-2: Mean values of number of viable colonies of bacteria in each group at 10^{-3} dilution

Group	Number viable colonies (MEAN\pmSD)
Group-I	578.39 \pm 1.67
Group-II	0.00 \pm 0.00*
Group-III	0.00 \pm 0.00*
Group-IV	0.00 \pm 0.00*
Group-V	0.00 \pm 0.00*

Table-3: Mean values of number of viable colonies of bacteria in each group at 10^{-6} dilution

Group	Number viable colonies (MEAN\pmSD)
Group-I	219.84 \pm 0.45
Group-II	0.00 \pm 0.00*
Group-III	0.00 \pm 0.00*
Group-IV	0.00 \pm 0.00*
Group-V	0.00 \pm 0.00*

Table-4: Multiple comparison of mean number of viable colonies of bacteria in the different groups

Group	Number viable colonies (MEAN\pmSD)
Group-I	1057.72 \pm 1.24
Group-II	0.00 \pm 0.00*
Group-III	0.00 \pm 0.00*
Group-IV	22.00 \pm 1.41* ^{#, \$}
Group-V	15.00 \pm 1.00* ^{#, \$, +}

Table-5: Mean values of number and percentage of samples with growth of bacteria in each group at 10^{-1} dilution

Group	Number of samples with growth of bacteria/Total number of samples	Percentage
Group-I	10/10	100%
Group-II	0/10	0% *
Group-III	0/10	0% *
Group-IV	9/10	90% *,#,†
Group-V	8/10	80% *,#,†,

Table-6: Mean values of number and percentage of samples with growth of bacteria in each group at 10^{-3} dilution

Group	Number of samples with growth of bacteria/total number of samples	Percentage
Group-I	10/10	100%
Group-II	0/10	0% *
Group-III	0/10	0% *
Group-IV	0/10	0% *
Group-V	0/10	0% *


Table-7: Mean values of number and percentage of samples with growth of bacteria in each group at 10^{-6} dilution

Group	Number of samples with growth of bacteria/total number of samples	Percentage
Group-I	10/10	100%
Group-II	0/10	0%*
Group-III	0/10	0%*
Group-IV	0/10	0%*
Group-V	0/10	0%*


Table-8: Multiple comparisons of mean values of number and percentage of samples with growth of bacteria in between groups

Group	Number of samples with growth of bacteria/total number of samples	Percentage
Group-I	10/10	100%
Group-II	0/10	0%*
Group-III	0/10	0%*
Group-IV	9/10	90%*,#,\$
Group-V	8/10	80%*,#,\$,†

Graph-1: Mean values of number and percentage of samples with growth of bacteria in each group at 10^{-1} dilution



Graph-2: Multiple comparison of mean number of viable colonies of bacteria in the different groups



Graph-3: Multiple comparisons of mean values of number and percentage of samples with growth of bacteria in between groups






Fig 1: Specimens stored in normal saline after biomechanical preparation



Fig 2: Specimens immersed in streptococcus selection broth for autoclaving



Fig 3: Specimens subjected to autoclaving




Fig 4: Laminar flow



Fig 5: Inoculation of specimens with *E.faecalis* suspension using tuberculin syringes.



Fig 6: Specimens kept for incubation




Fig 7: SEM picture showing *E. faecalis* colonies



Fig 8: Irrigation of specimens with the test irrigants



Fig 9: Sample preparation using H files.



Fig 10: Collecting dentinal shavings in test tube containing normal saline





Fig 11: Agar plates showing complete inhibition by 5.25% NaOCl



Fig 12: Agar plate showing complete inhibition by 2% CHX

Fig 13: Agar plate (Garlic) showing a) complete inhibition b) presence of CFU

A



B






Fig 14: Agar plate (Cinnamon) showing a) complete inhibition b) presence of CFU

A



B






Fig 15: Agar plates (saline) showing innumerable CFU

Bibliography

1. Kakehashi S, Stanley H R, Fitzgerald R J. The effects of surgical exposure of dental pulps in germ-free and conventional laboratory rats. *Oral surgery, Oral medicine and oral pathology* 1965;Vol 20:34-5.
2. Sundqvist G. Ecology of the root canal flora .*J Endod* 1992;Vol 18:142-149
3. Sjogren U, Figdor D, Spangberg L, Sundqvist G. Influence of infection at the time of root filling on the outcome of endodontic treatment of teeth with apical periodontitis. *International Endodontic Journal* 1997; Vol 30:297-306.
4. Bystrom A, Sundqvist G. Bacteriologic evaluation of 0.5% NaOCl in endodontic therapy. *Oral surgery, Oral medicine and Oral pathology* 1983; Vol 55:307-312.
5. Haapasalo M,Orstavik D.In vitro infection and disinfection of dentinal tubules. *Journal of Dental Research* 1987; Vol 66:1375-1379.
6. Engstrom B. The Significance of Enterococci in Root Canal Treatment. *Odontologisk Revy* 1964; Vol15: 87-106.
7. R.M. Love. *Enterococcus faecalis- A mechanism for its role in endodontic failure. International Endodontic Journal* 2001; Vol 34: 399-405.
8. Sebeena Mathew, Boopathy. *Enterococcus faecalis - An endodontic challenge. JIADS* 2010; Vol 1(4):33-37
9. Isabelle Portenier, Tuomnos M T, Waltimo, Markus Haapasalo. *Enterococcus Faecalis – the root canal survivor and ‘star’ in post – treatment disease. Endodontic Topics* 2003; Vol 6:135-159.

10. Sushma Jaju, Prashant P.Jaju. Newer Root Canal Irrigants In Horizon: A Review. *International Journal of Dentistry* 2011;1-9
11. Matthias Zehnder. Root Canal Irrigants. *J Endod* 2006; Vol 32 (5): 389 - 398.
12. Deivanayagam Kandaswamy, Nagendrababu Venkatesh Babu. Root canal Irrigants. *Journal of Conservative Dentistry* 2010; Vol 13(4):256-264.
13. Madhu Pujar, ChetanPatil, AjayKadam. Comparison of antimicrobial efficacy of Triphala, Green tea polyphenols and 3% Sodiumhypochlorite on *Enterococcus faecalis* biofilm formed on tooth substrate – In vitro. *JIOH* 2011; Vol 3 (2) : 23-29.
14. Zahed Mohammadi. An update on the antibiotic –based root canal irrigation solutions. *International Endodontic Journal* 2008; Vol 3(2): 1-7
15. Alamsher. Antimicrobial activity of natural products from medicinal plants. *Gomal Journal of Medical Science* 2009; Vol 7 (1):72-78.
16. Meenal. N Gulve, Nitin.D. Gulve. Comparison of antimicrobial efficacy of Ginger extract and 2%Sodium hypochlorite against *Enterococcus faecalis* using Agar diffusion method .*JIDA* 2010; Vol 4(10):347-349.
17. MadhuPujar, SaleemMakandar. Herbal usage in endodontics A Review. *Int. Journal of Contemporary Dentistry* 2011; Vol 2 (1) :34-37.
18. Ramkumar Pundir, Pranay Jain, Chetan Sharma. Antimicrobial activity of ethanolic extracts of *Syzygium Aromaticum* and *Allium Sativum* against food associated bacteria and fungi. *Ethnobotanical Leaflets* 2010; Vol 14:344-360.

19. M.D MahfuzulHoque, M.L Bari, Vijay K Juneja, S. Kawamoto. Antimicrobial activity of cloves and cinnamon extracts against food borne pathogens and spoilage bacteria and inactivation of *Listeria Monocytogenes* in ground chicken meat with their essential oils. Rep. Nat' I. Food Res. Inst 2008; Vol 72:9-21.
20. SrinivasanDurairaj, Sangeetha Srinivasan, P.Lakshmanaperumalsamy. Invitro antibacterial activity and stability of Garlic extract at different PH and temperature. Electronic Journal of Biology 2009; Vol5 (1): 5-10.
21. Dhinahar. S, Lakshmi. T. Role of botanicals as antimicrobial agent in management of dental infections – A Review. International Journal of Pharma And Biosciences 2011; Vol2(4): 690-704.
22. Enzo A. Palombo. Traditional medicinal plant extract and natural products with activity against oral bacteria. Potential application in the prevention and treatment of oral Diseases. Evidence Based Complementary and Alternative Medicine 2011 : 1-12.
23. Dhanyakumar,N M PreenaSidhu. The antimicrobial activity of *AzadirachtaIndica*, *Glycyrrhizaglabra*, *Cinnamumzeylanicum*, *SyngiumAromaticum*, *AccaciaNilotica* on *streptococcus Mutans* and *Enterococcus Faecalis*-An in vitro study.Endodontology:18-25.
24. R.R. White, G.L. Hays, L.R. Janer. Residual antimicrobial activity after canal irrigation with Chlorhexidine. JEndod 1997; Vol 23(4):229-231.
25. L.B. Peters, P.R. Wesselink, W.R. Moorer. Penetration of bacteria in bovine root dentine In Vitro. International Endodontic Journal 2000; Vol 33: 28-36.

26. Shyh-mingTsao,Mei-chin Yin. In vitro activity of Garlic oil and four diallylsulphides against antibiotic resistant *Pseudomonas aeruginosa* and *Klebsiella pneumonia*. *Journal of Antimicrobial Chemotherapy*2001; Vol 47:665-670.
27. B.P.F.A. Gomes, C.C.R. Ferraz, M.E.Vianna, V.B.Berber, F.B.Teixeira, F.J.Souza-Filho. In Vitro antimicrobial activity of several concentrations of sodiumhypochlorite and chlorhexidinegluconate in the elimination of *Enterococcus faecalis*. *International Endodontic Journal* 2001; Vol 34: 434-438.
28. Bettina Basrani, J.Miguel Santos, Leo Tjaderhane, Helen Grad, Omer Gorduysns, Junfu Huang, Herenia.P. Lawrence, Shimon Friedman. Substantive antimicrobial activity in Chlorhexidine-treated human root dentin. *Oral Surg, Oral Med, Oral Pathol, Oral Radiol,Endod* 2002; Vol 94: 240-245.
29. J.C. Yamashita, M. Tanomaru Filho, M.R. Leonardo, M.A. Rossi L.A.B. Silva. Scanning Electron Microscopic study of the cleaning ability of Chlorhexidine as a root canal irrigant. *International Endodontic Journal* 2003; Vol 36: 391-394.
30. Onyeagba R.A, Ugbogu O.C,Okeke. C.U,Iroakasi O. Studies on the antimicrobial effects of garlic, ginger and lime. *African Journal of Biotechnology* 2004; Vol 3(10) : 552-554.
31. I.M. Saleh, I.E. Ruyter, M. Haapasalo, D. Orstavik. Survival of *Enterococcus faecalis* in infected dentinal tubules after root canal filling with different Root Canal Sealers In Vitro. *International Endodontic Journal* 2004; Vol37 : 193-198.

32. C.E. Radcliffe, L. Potouridou, R. Qureshi, N. Habahbeh, A. Qualtrough, H. Worthington, D. B. Drucker. Antimicrobial activity of varying concentrations of Sodium hypochlorite on the endodontic microorganisms *Actinomyces israelii*, *A. naeslundii*, *Candida albicans* and *Enterococcus faecalis*. *International Endodontic Journal* 2004; Vol 37: 438-446.
33. Katherine R. Carson, Gary G. Goodell, Scott B, McClanaban. Comparison of the antimicrobial activity of six irrigants on primary endodontic pathogens. *J Endod* 2005; Vol 31(6):471-473.
34. Fabio Roberto Dametto, Caio Cezar Randi Ferraz, Brenda Paula, Figueiredo de Almeida Gomes, Alexandre Augusto Zaia, Fabricio Batista Teixeira, Francisco Jose de Souza-Filho, Piracicaba. In vitro assessment of the immediate and prolonged antimicrobial action of chlorhexidine gel as an endodontic irrigant against *Enterococcus faecalis*. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2005; Vol 99 F68-772.
35. Brain D. Barnhart, Augustine Chuang, Jurandir J. Daile Lucca, Steven Roberts, Frederick Liewehr, Anthony. P. Joyce. An In vitro evaluation of the cytotoxicity of various endodontic irrigants on human gingival fibroblast. *J Endod* 2005; Vol 31(8) : 613-615
36. Russell. S. Eddy, Anthony P. Joyce, Steve Roberts, Thomas. B. Buxton, Frederick Liewehr. An In vitro evaluation of the antibacterial efficacy of Chlorine Dioxide on *E. Faecalis* in bovine Incisors – *J Endod* 2005; Vol 31 (9) : 672-675.

37. Suree Nanasombat, Pana Lohassupthawee. Antibacterial activity of crude ethanolic extracts and essential oils of spices against Salmonellae and other Enterobacteria. KMITL Sci. Tech. J 2005; Vol 5(3): 527-538.
38. Melissa L. Ruff, Scott B. McClanahan, Britta S. Babel. In Vitro antifungal efficacy of four irrigant as a final rinse. J Endod 2006;Vol 32 (4):331-333.
39. M.S. Clegg, F.J Vertucci, C. Walker, M. Belanger, L.R. Britto. The effect of exposure to irrigant solutions on apical dentin biofilm in vitro. J. Endod 2006; Vol 32 (5): 434-437.
40. Thomas R. Dunavant, John. D. Regan, Gerald N. Glickman, Eric S. Solomon, Allen L. Honeyman. Comparative evaluation of endodontic irrigants against Enterococcusfaecalis Bio Filims. J Endod 2006; Vol 32 (6): 527 – 531.
41. Isabelle Portenier, TuomasWaltimo, Dag Orstavik, Markus Haapasalo. Killing of Enterococcus faecalis by MTAD AndChlorhexidineDigluconate with or without Cetrimide in the presence or absence of dentin Powder or BSA. J Endod 2006; Vol 32 (2) :138 – 141.
42. V.B Berber, B.P.F.A Gomes. N. T. Sena, C.C.R. Ferraz, A.A. Zaia, F. J. Souza–Filho. Efficacy of various concentrations of NaOCl and instrumentation techniques in reducing Enterococcus faecalis within root canals and dentinal tubules. International Endodontic Journal 2006, Vol 39: 10-17.
43. N.T Sena, B.P.F.A Gomes, M.E Vianaa, V.B Berber, A.A Zaia,C.C. R Ferraz, F.J Souza–Filho. In vitro antimicrobial activity of sodium

- hypochlorite and chlorhexidine against selected single-species biofilm.
International Endodontic Journal 2006;Vol 39(11):878-885.
44. Seung – Eun Yang, Jeong – Heon cha, Eui-Seong Kim, Kee – YeonKum, Chang – Young Lee, H-Young Jung. Effect of smearlayer and Chlorhexidine treatment on the adhesion of *Enterococcus faecalis* to bovine dentin. J Endod 2006; Vol 32 (7); 663 – 667.
45. John.M. Williams, Martin Trope, Daniel. J. Caplan, Diane C. Shugars. Detection And quantitation of *Enterococcus faecalis* by Real – time PCR (QPCR), Reverse Transcription PCR (RT-PCR) and cultivation during endodontic treatment. J Endod; Vol 32 (8): 715 – 721.
46. Erika M. Johnson, Susan E. Flannagan, Christine M. Sedgley. Coaggregation interactions between oral and endodontic *Enterococcus faecalis* and bacterial species isolated from persistent apical periodontitis. J Endod 2006; Vol 32 (10); 946 – 950.
47. Bettina. R. Basrani, Sheela Manek, Rana N.S.Sodhi, Edward Fillery, Aldo Mazur. Interaction between Sodium hypochlorite and chlorhexidine Gluconate. J Endod 2007; Vol 33(8): 966-969.
48. Bradley. M. Newberry, Shahrokh Shabahang, Neal Johnson, Raydolfo. M. Aprecio, Mahmoud Torabinejad. The antimicrobial effect of biopure MTAD on eight strains of *Enterococcus faecalis*: An In vitro investigation. J Endod 2007; Vol 33(11); 1352-1354.
49. Trisha. A. Krause, Frederick. R. Liewehr, Chin-Lo Hahn. The antimicrobial effect of MTAD, Sodium hypochlorite, Doxycycline and Citric Acid on *Enterococcus Faecalis*. J Endod 2007; Vol33 (1): 28-30.

50. Luciano Giardino, Emanule Ambu, Enrico Savoldi, Roberto Rimondini, Clara Cassanelli, Eugenio A. Debbia. Comparative evaluation of antimicrobial efficacy of Sodium Hypochlorite, MTAD, and Tetraclean against *Enterococcus faecalis* biofilm. *J Endod* 2007; Vol20(10): 1-4.
51. Caio.C.R. Ferraz, Brenda. P.F.A. Gomes, Alexandre .A. Zaia, Fabricio. B. Teixeira, Francisco. J.Souza-Filho. Comparative study of the antimicrobial efficacy of Chlorhexidine Gel, Chlorhexidine Solution and Sodium hypochlorite as endodontic irrigants. *Braz Dent J* 2007; Vol 18(4):294-298.
52. J.Craig Baumgartner, Stephen Johal, J. Gordon Marshall. Comparison of the antimicrobial efficacy of 1.3% NaOCl/Bio pure MTAD to 5.25% NaOCl/15% EDTA for Root canal Irrigation. *J Endod* 2007; Vol33 (1) : 48-51.
53. Mathew J. Royal, Anne. E.Williamson, David. R. Drake. Comparison of 5.25% NaOCl, MTAD and 2% Chlorhexidine in the rapid disinfection of Polycaprolactone based root canal filling material. *J Endod* 2007; Vol33 (1) : 42-44.
54. Monika Marending, Frank Paque, Jens Fischer, MalthiasZehnder. Impact of irrigants sequence on mechanical properties of human root dentin. *J Endod* 2007; Vol33(11): 1325-1328
55. Jose. F. Siqueira, Simone S.M.Paiva, Isabela. N. Rocas. Reduction in the cultivable bacterial populations in infected root canals by a Chlorhexidine-Based antimicrobial protocol. *J Endod* 2007; Vol 33(5): 541-547.

56. William. J. Nudera, Mohamed. I. Fayad, Bradford. R. Johnson, Min Zhu, Christopher. S.Wenckus, Ellen. A. Begole, Christine D. Wu. Antimicrobial effect of Triclosan and Triclosan with Gantrez on five common endodontic pathogens. JEndod 2007; Vol 35(10):1239-1242.
57. Joshua M. Davis, James Maki, James K Bahcall. An invitro comparison of the antimicrobial effects of various endodontic medicaments on E.faecalis. J Endod 2007; Vol 33(5):567-569.
58. Sunitha Bansod, Mahendra Rai. Antifungal activity of essential oils from Indian Medicinal plants against human pathogenic *Aspergillus fumigatus* and *A. Niger*. World journal of Medical Sciences 2008; Vol 3 (2): 81 – 86.
59. Zohresh Ahangari, Mohammad Samiee, Mohammad Amin Yolmeh, Gita Eslami. Antimicrobial activity of three root canal irrigants on *Enterococcus Faecalis*. An In vitro study. International Endodontic Journal 2008; Vol 3 (2) : 33-37.
60. P. Chivatxaranukul, S.G. Dashaper, H.H Messer. Dentinal tubule invasion and adherence by *Enterococcus Faecalis*. International Endodontic Journal 2008; Vol 41: 873 – 882.
61. Rachele Neglia, Andrea Ardizzoni, Luciano Giardino, Emmanuele Ambu, Silvia Grazi, Sara Calignano, Cinzia Rimoldi, Elena Righi, Elisabetta Blasi. Comparative in vitro and ex vivo studies on the bactericidal activity of Tetraclean, A new generation endodontic irrigant and Sodium Hypochlorite. New Micro Biologica 2008; Vol 31: 57-65.

62. Sandeep Singh, Sushma Krishna, Shashi Rashmi Aharya, P.Sugandhi Rao, Vasudev Ballal. Comparison of antimicrobial efficacy of Biopure MTAD and 2.5% Sodium Hypochlorite irrigation in infected root canals following single visit endodontics – An In vivo study. *Endodontology* 2009; Vol 21 (1): 27-33.
63. S. Shobhana, V.G Vidhya, M. Ramya. Antibacterial activity of Garlic varieties (*Ophioscordon* and *Sativum*) on enteric pathogens. *Current Research Journal of Biological Sciences* 2009; Vol 1 (3) :123 – 126.
64. Andrea Ardizzoni, Elisabeth Blasi, Cinzia Rimoldi, Luciano Giardino, Emmanuele Ambu, Elena Righi, Rachele Neglia. An In vitro and Ex vivo study on two antibiotic-based Endodontic Irrigants- A challenge To Sodium hypochlorite. *New Microbiologica* 2009; Vol 32 : 57-66.
65. YaShen ,WeiQian , Christopher Chung, Ingar Olsen, Markus Haapasalo. Evaluation of the effect of two chlorhexidine preparations on biofilm bacteria In vitro: A three –dimensional quantitative analysis. *J Endod* 2009; Vol 35(7):981-985.
66. Rosina Khan, Barira Islam, MohdAkram, Shazi Shakil, Anis Ahmad, S. Manazir Ali, Mashiatullah Siddiqui, Azad.U.Khan. Antimicrobial activity of five herbal extracts against multi drug resistant strains of bacteria and fungus of clinical origin. *Molecules* 2009; Vol 14: 586-597.
67. Bishnu Joshi, Sunil Lekhak, Anuja Sharma. Antibacterial property of different medicinal plants: *Ocimum Sanctum*, *Cinnamomum Zeylanicum*, *Xanthoxylum Armatum* and *Origanum Majorana*. *Kathmandu University Journal Of Science, Engineering And Technology* 2009; Vol5 (1):143-150.

68. Luciano Giardino, Enrico Savoidi, Emmanuel Ambu, Roberto Rimondini, Alberto Palezona, Eugennio.A.Debbia. Antimicrobial effect of MTAD, Tetraclean, Cloreximid and Sodium Hypochlorite on three common Endodontic Pathogens. *Indian J Dent Res* 2009; Vol20(3):12-16.
69. U.B.Owhe-Urghe, D.A. Ehwarieme, D.O.Eboh. Antibacterial activity of Garlic and Lime on isolates of extracted carious Teeth. *African Journal of Biotechnology* 2010;Vol 9(21): 3163-3166.
70. Vahid Zand, Mehrdad Lofti, Saeed Rahimi, Hadi Mokhtari Ali Kazemi, Vahideh Sakhamanesh. A comparative Scanning Electron Microscopic investigation of the smear layer after the use of Sodium hypochlorite Gel and solution form as root canal irrigants. *J Endod* 2010; Vol 36(7); 1234-1237.
71. Carrmen Maria Ferrer-Luque, Maria Teresa Arias-Moliz, Maria Paloma Gonzalez-Rodriguez, Pilar Baca. Antimicrobial activity of Maleic acid and combinations of cetrimide with chelating agents against *Enterococcus faecalis* biofilm. *J Endod* 2010; vol 36(10): 1673-1675.
72. Saurabh. S. Chandra, Revathi Miglani, M.R. Srinivasan, Rajamani Indira, Antifungal efficacy of 5.25% Sodium Hypochlorite, 2% Chlorhexidine Gluconate, and 17% EDTA with and without an antifungal Agent. *J Endod* 2010; Vol 36(4):675-678.
73. Andrea Cruz Camara, Miracy Muniz De Albuquerque, Carlos MenzesAguiar, Ana Cristina Regis De Barros Correia. Antimicrobial activity of Chlorhexidine in root canals instrumented with the Protaper universal System. *Braz Joral Sci* 2010; Vol 9(3): 402-409.

74. Maria Teresa Arias-Moliz, Carmen Maria Ferrer-Luque, Maria Paloma Gonzalez-Rodriguez, Mariano Jose Valderrama. Eradication of *Enterococcus faecalis* biofilm by Cetrimide and Chlorhexidine. *J Endod* 2010; Vol 36(1) : 87 -90.
75. Luis F. Chavez de Paz, Gunnar Bergenholtz, Gunnel S Vensater. The effects of antimicrobials on endodontic biofilm bacteria. *J Endod* 2010; Vol 36 (1) : 70 – 77.
76. Bonnic Retamozo, Shahrokh Shabahang, Neal Johnson Raydolfo. M.Aprecio, Mahmoud Torabinejad. Minimum contact time and concentration of Sodium Hypochlorite required to eliminate *Enterococcus faecalis*. *J Endod* 2010, Vol 36(3): 520-523.
77. J.Prabhakar, M.Senthilkumar, M.S.Priya, K. Mahalakshmi, P.K. Sehgal, V.G. Sukumaran. Evaluation of antimicrobial efficacy of herbal alternatives (Triphala and Green Tea Polyphenols), MTAD and 5% Sodium Hypochlorite against *Enterococcus faecalis* biofilm formed on tooth substrate. An in vitro study. *J Endod* 2010; vol36 (1): 83-86.
78. F.G. Pappen, Y. Shen, W.Aian, M. R. Leonardo, L.Giardino, M.Haapasalo. In vitro antibacterial action of Tetraclean, MTAD and five experimental irrigation solutions. *International Endodontic Journal*; Vol 43:528-535.
79. Sonja Stojicic, Slavoljub Ziukovic, Wei Qian, Huizhang, Markus Haapasalo. Tissue Dissolution by Sodium Hypochlorite. Effect of Concentration, Temperature, Agitation and Surfactant. *J Endod* 2010; Vol 36(9): 1558-1562.

80. Isabela N. Rocas, Jose. F. Siqueira. Identification of bacterial enduring endodontic treatment procedures by a combined Reverse Transcriptase – Polymerase Chain Reaction and Reverse- Capture Checker board Approach. J Endod 2010; Vol 36(1) : 45-52.
81. Moeen Mahmoud Al Weshah. The in vitro effect of 2% chlorhexidine on dentin hardness. Pakistan Oral and Dental Journal 2011; Vol 31 (1) : 173-177.
82. Zahed Mohammadi, Ali Mombeinipour, Luciano Giardino, Shahriar Shaahriari. Residual antibacterial activity of a new modified Sodium Hypochlorite - based endodontic irrigation solution. Med Oralpatol Oralcir Bucal 2011; Vol 16 (4) : 588 – 592
83. Poonam Shingare, Vishwas Chaugule. Comparative evaluation of antimicrobial activity of Miswak, Propolis, Sodium Hypochlorite and Saline as root canal irrigants by microbial culturing and quantification inchronically exposed primary teeth. GERMS 2011; Vol 1 (1): 12-21.
84. AshishSaraf, Mohit. S Mishra, K. Sharama. Antibacterial activity of commercial and wild Cinnamon species. Journal of Phytology 2011; Vol 3 (2): 102 – 106.
85. Zohreh Dalirsani, Mohammad Adibpour, Mohammad Aghazadeh Maryam Amirchagmaghi, Farnaz Falaki, Pegah Mosannen, Faezeh Mojarad Hamzri. Invitro Comparison of inhibitory activity of 10 plant extract against Candida albicans. Australian Journal of basic and Applied Sciences 2011;Vol 5 (5) : 930 – 935.
86. Mehrdad Lotfi, Sepideh Vosoughhosseini, Bahram Ranjkesh, Sajjadkhani, Mohammadali Saghiri, Vahid Zand. Antimicrobial efficacy

- of Nanosilver, Sodium Hypochlorite and Chlorhexidinegluconate against *Enterococcus faecalis*. *African Journal of Biotechnology* 2011; Vol 10 (35): 6799 – 6803.
87. Melahat Gorduysus, Behram Tuncel, Emre Nagas, Koray Grgunay, Pinar Yurdakul, Sibel Erguven, Ozgur Yildirim Torun, Omer Gorduysus. Antimicrobial effects of various endodontic irrigants on selected microorganisms. *Clinical Dentistry and Research* 2011; Vol 35 (1): 41 - 46.
88. M.N Shahni, V. V. Subba Reddy. Comparison of antimicrobial substantivity of root canal irrigants in instrumented root canal up to 72 hrs. An in vitro study. *Journal of Indian Society of Pedodontics And Preventive Dentistry* 2011; Vol 29 (1) : 28-33.
89. Ponmurugan Karuppiah, Shyamkumar Rajaram. Antibacterial effect of *Allium Sativum*, cloves And *Zingiber Officinale* rhizome against multiple drug resistant clinical pathogens. *Asian pacific Journal of Tropical Biomedicine* 2012: 597 – 601.
90. D. Mortenson, M. Sadilek, N.M. Flake, A Paranjpe, I Heling, J.D Johnson. The effect of using an alternative irrigant between Sodium Hypochlorite and Chlorhexidine to prevent the formation of Para – Chloroaniline within the root canal system. *International Endodontic Journal* 2012: 1-5.
91. S. Stojicic, Y. Shen, W. Qian, B. Johnson, M. Haapasalo. Antibacterial and smear layer removal ability of a novel irrigant, Q Mix. *International Endodontic Journal* 2012; Vol 45 : 363 – 371.

92. Abou-Rass M, Piccinino MV. The effectiveness of four clinical irrigation methods on the removal of root canal debris. Oral surgery, Oral medicine and Oral pathology 1982;Vol54:323-328.
93. Lynne RE, Lieweher FR, Patton WR, Buxton TB, McPherson JC. In Vitro Antimicrobial Activity of Various Medication Preparations on E.faecalis in Root Canal Dentin .J Endod 2003;Vol29:187-190.
94. Orstavik D, Haapasalo M. Disinfection by endodontic irrigants and dressings of experimentally infected dentinal tubules. Endodontics and Dental Traumatology 1990;Vol6:142-149.
95. Hancock HH, Sigurdsson A, Trope M, Moiseiwitsch J. Bacteria isolated after unsuccessful endodontic treatment in a North American population. Oral surgery, Oral medicine, Oral pathology, Oral radiology and Endodontics 2001;Vol91:579-586.
96. Siqueira JF Jr, Machado AG, Silveria RM, Lopes HP, Uzeda M. Evaluation of the effectiveness of sodium Hypochlorite used with three irrigation methods in the elimination of E.faecalis from root canals in vitro. International Endodontic Journal 1997;Vol30:279-282.
97. Akpata ES, Blechman H. Bacterial invasion of pulpal dentine wall in vitro. Journal of Dental Research 1982;Vol61:435-438.
98. Khademi A, Yazdizadeh M, Feizianfard M. Determination of the minimum instrumentation size for penetration of irrigants to the apical third of root canal systems. J Endod 2006;Vol32:417-420.
99. Safavi KE, Spanberg L, Langeland K. Smear layer removal effects on root canal dentin tubule disinfection. J Endod 1989;Vol15:175-176.

100. Adriaens PA, de Boever JA, Loesche WJ. Bacterial invasion in root cementum and radicular dentine of periodontally teeth in humans. *Journal of Periodontology* 1988;Vol59;222-230.
101. Siqueira JF, de Uzeda M, Fonseca MEF. A scanning electron microscopic evaluation of in vitro dentinal tubule penetration by selected anaerobic bacteria. *J Endod* 1996;Vol22:308-310.
102. Eldeniz AU, Ozer F, Hadimli HH, Erganis O. Bactericidal activity of Er,Cr:YSGG laser irradiation against *E. faecalis* compared with NaOCl irrigation: an ex vivo pilot study. *International Endodontic Journal* 2007;Vol40(2):112-9.
103. Siqueira JF, Rocas IN, Faviera A, Lima KC. Chemomechanical reduction of the bacterial population in the root canal after instrumentation and irrigation with 1.0, 2.5 and 5.25% NaOCl. *J Endod* 2000;Vol26:331-4.
104. Vianna ME, Gomes BP, Berber VB, Zaia AA, Ferraz CC, de Souza-Filho FJ. In vitro evaluation of the antimicrobial activity of Chlorhexidine and Sodium Hypochlorite. *Oral Surg, Oral Med, Oral Pathol, Oral Radiol, Endod* 2004;Vol97:79-84.
105. H. K. Haapasalo, E. K. Siren, T M T Waltimo, D .Orstavik, M.P.P. Haapasalo. Inactivation of local root canal medicaments by dentin: an in vitro study. *International Journal of Endodontics* 2000; Vol33:126-131.
106. Spangberg L, Engstrom B, Langeland K. Biological effects of dental materials. Toxicity and antimicrobial effect of endodontic antiseptics in vitro. *Oral Surg, Oral Medicine and Oral Pathology* 1973;Vol36:856-870.

107. Fardal O, Turnbull RS. A review of the literature on use of chlorhexidine in dentistry. *Journal of the American dental association* 1986; Vol 112:863-869.
108. Ohara PK, Torabinejad M, Kettering JD. Antibacterial effects of various endodontic irrigants on selected anaerobic bacteria. *Endodontics and Dental Traumatology* 1993; Vol 9:95-100.
109. Oncag O, Hosgor M, Hilmioglu S, Zekioglu O, Eronat C, Burhanoglu D. Comparison of antibacterial and toxic effects of various root canal irrigants. *International Endodontic Journal* 2003; Vol 36:423-32.
110. Kuruvilla JR, Kamath MP. Antimicrobial activity of 2.5% sodium Hypochlorite, 0.2% chlorhexidine gluconate separately and combined as endodontic irrigants. *J Endod* 1998; Vol 24:472-6.
111. Rosenthal S, Spanberg I, Safavi KE. Chlorhexidine substantivity in root canal dentine. *Oral Surg, Oral Med, Oral Pathol, Oral Radiol, Endod* 2004; Vol 98:488-92.
112. Ekta Menghani, Arvind Pareek, R.S.Negi, C K Ojha. Search for antimicrobial potentials from certain Indian medicinal plants. *Research Journal of Medicinal Plant* 2011; Vol 5(3):295-301.
113. Sana Mukhtar, IfraGhori. Antibacterial activity of aqueous and ethanolic extracts of garlic, cinnamon and turmeric against *Eschereshia coli* ATCC25922 and *Bacillus Subtilis* DSM 3256. *International journal of Applied Biology and Pharmaceutical Technology* 2012; Vol 3(2):131-136.
114. Breed R, W D Dotterer. The number of colonies allowable on satisfactory agar plates. *J Bacteriol* 1916; Vol 1:321-33.

115. B.E.Boboye, A.J.Alli. Cellular effects of Garlic extract on *Pseudomonas aeruginosa* and *Staphylococcus aureus*. *Research Journal of Medicinal Plant*2008;Vol2(2):79-85.
116. Clem Maidment, ZosiaDembny, Paul King. Investigations into the antibacterial properties of garlic using the disc assay method. *Journal of Biological Education*1998;Vol32(3):162-165.